




Rec'd PCT/AT 25 MAY 2005
PCT/GB 2003 / 0 0 4 6 3 1

27 OCTOBER 2003

INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ




I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *Andrew Jones*

Dated 19 September 2003

**PRIORITY
DOCUMENT**

BEST AVAILABLE COPY SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



27NOV02 E76610-3 002093
P01/7700 0.00-0227555.0

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference
PPD 70125/GB/P

2. Patent application number
(The Patent Office will fill in this part)
26 NOV 2002 0227555.0

3. Full name, address and postcode of the or of each applicant (underline all surnames)
SYNGENTA Limited
European Regional Centre
Priestley Road
Surrey Research Park, Guildford,
Surrey, GU2 7YH, United Kingdom

Patents ADP number (if you know it)

6254007002

If the applicant is a corporate body, give the country/state of its incorporation

UNITED KINGDOM

833074800

4. Title of the invention
FUNGICIDES

5. Name of your agent (if you have one)
Malcolm John HOUGHTON
Intellectual Property Department
Syngenta Limited
Jealott's Hill International Research Centre
PO Box 3538
Bracknell, Berkshire, RG42 6YA
UNITED KINGDOM

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Patents ADP number (if you know it)

8157760001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body.

See note (d))

YES (b)

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form	50
Description	03
Claim(s)	00
Abstract	00
Drawing(s)	00

10. If you are also filing any of the following, state how many against each item.

Priority documents	
Translations of priority documents	
Statement of inventorship and right to grant of a patent (<i>Patents Form 7/77</i>)	
Request for preliminary examination and search (<i>Patents Form 9/77</i>)	
Request for substantive examination (<i>Patents Form 10/77</i>)	
Any other documents (<i>please specify</i>)	

11.

I/We request the grant of a patent on the basis of this application.
Syngenta Limited

Signature *J A Bardich*
Authorised Signatory

Date *26 / 11 / 02*

12. Name and daytime telephone number of person to contact in the United Kingdom

Joanna Carmen CHANDLER = 01344 414365

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes


- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*
- For details of the fee and ways to pay please contact the Patent Office.*

FUNGICIDES

This invention relates to novel *N*-alkynyl-2-quinolin-(isoquinolin- and quinazolin-)oxyalkylamides, to processes for preparing them, to compositions containing
 5 them and to methods of using them to combat fungi, especially fungal infections of plants.

Various quinolin-8-oxyalkanecarboxylic acid derivatives are described as being useful as antidotes for herbicides or as herbicide safeners (see, for example, US 4,881,966, US 4,902,340 and US 5,380,852). Certain pyridyl- and pyrimidinyl-
 10 (thio)alkanoic acid amide derivatives are described in, for example, WO 99/33810 and US 6090815, together with their use as agricultural and horticultural fungicides. In addition, certain phenoxyalkanoic acid amide derivatives are described in, for example, US 4116677 and US 4168319, together with their use as herbicides and mildewicides.

According to the present invention, there is provided a compound of the general
 15 formula (1):



wherein one of X and Y is N and the other is CR or both of X and Y are N; R is H, halo, C₁₋₈ alkyl, C₃₋₆ cycloalkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, C₁₋₈ alkoxy, C₁₋₈ alkylthio, nitro,
 20 amino, mono- or di-(C₁₋₆)alkylamino, mono- or di-(C₂₋₆)alkenylamino, mono- or di-(C₂₋₆)alkynylamino, formylamino, C₁₋₄ alkyl(formyl)amino, C₁₋₄ alkylcarbonylamino, C₁₋₄ alkyl(C₁₋₄ alkylcarbonyl)amino, cyano, formyl, C₁₋₄ alkylcarbonyl, C₁₋₄ alkoxy carbonyl, aminocarbonyl, mono- or di-(C₁₋₄)alkylaminocarbonyl, carboxy, C₁₋₄ alkylcarbonyloxy, aryl(C₁₋₄)alkylcarbonyloxy, C₁₋₄ alkylsulphinyl, C₁₋₄ alkylsulphonyl or C₁₋₄ alkyl-
 25 sulphonyloxy; R₁ is C₁₋₄ alkyl, C₂₋₄ alkenyl or C₂₋₄ alkynyl in which the alkyl, alkenyl and alkynyl groups are optionally substituted on their terminal carbon atom with one, two or three halogen atoms (e.g. 2,2,2-trifluoroethyl), with a cyano group (e.g. cyanomethyl), with a C₁₋₄ alkylcarbonyl group (e.g. acetylmethyl), with a C₁₋₄ alkoxy carbonyl group (e.g. methoxycarbonylmethyl and methoxycarbonylethyl) or with a hydroxy group (e.g.

hydroxymethyl), or R₁ is alkoxyalkyl, alkylthioalkyl, alkylsulphinyalkyl or alkylsulphonylalkyl in which the total number of carbon atoms is 2 or 3 (e.g. methoxymethyl, methylthiomethyl, ethoxymethyl, 2-methoxyethyl and 2-methylthioethyl), or R₁ is a straight-chain C₁₋₄ alkoxy group (i.e. methoxy, ethoxy, *n*-propoxy and *n*-butoxy); R₂ is H, C₁₋₄ alkyl, C₁₋₄ alkoxymethyl or benzyloxymethyl in which the phenyl ring of the benzyl moiety is optionally substituted with C₁₋₄ alkoxy; R₃ and R₄ are independently H, C₁₋₃ alkyl, C₂₋₃ alkenyl or C₂₋₃ alkynyl provided that both are not H and that when both are other than H their combined total of carbon atoms does not exceed 4, or R₃ and R₄ join with the carbon atom to which they are attached to form a 3 or 4 membered carbocyclic ring optionally containing one O, S or N atom and optionally substituted with halo or C₁₋₄ alkyl; and R₅ is H, C₁₋₄ alkyl or C₃₋₆ cycloalkyl in which the alkyl or cycloalkyl group is optionally substituted with halo, hydroxy, C₁₋₆ alkoxy, C₁₋₆ alkylthio, cyano, C₁₋₄ alkylcarbonyloxy, aminocarbonyloxy or mono- or di(C₁₋₄)alkylaminocarbonyloxy, tri(C₁₋₄)alkylsilyloxy, optionally substituted phenoxy, optionally substituted thienyloxy, optionally substituted benzyloxy or optionally substituted thienylmethoxy, or R₅ is optionally substituted phenyl, optionally substituted thienyl or optionally substituted benzyl, in which the optionally substituted phenyl and thienyl rings of the R₅ values are optionally substituted with one, two or three substituents selected from halo, hydroxy, mercapto, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C₁₋₄ alkoxy, C₂₋₄ alkenyloxy, C₂₋₄ alkynyloxy, halo(C₁₋₄)alkyl, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, halo(C₁₋₄)alkylthio, hydroxy(C₁₋₄)alkyl, C₁₋₄ alkoxy(C₁₋₄)alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR'R'', -NHCOR', -NHCONR'R'', -CONR'R'', -SO₂R', -OSO₂R', -COR', -CR'=NR'' or -N=CR'R'', in which R' and R'' are independently hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C₁₋₄ alkyl or C₁₋₄ alkoxy.

The compounds of the invention contain at least one asymmetric carbon atom (and at least two when R₃ and R₄ are different) and may exist as enantiomers (or as pairs of diastereoisomers) or as mixtures of such. However, these mixtures may be separated into individual isomers or isomer pairs, and this invention embraces such isomers and

mixtures thereof in all proportions. It is to be expected that for any given compound, one isomer may be more fungicidally active than another.

Except where otherwise stated, alkyl groups and alkyl moieties of alkoxy, alkylthio, etc., suitably contain from 1 to 4 carbon atoms in the form of straight or
5 branched chains. Examples are methyl, ethyl, *n*- and *iso*-propyl and *n*-, *sec*-, *iso*- and *tert*-butyl. Where alkyl moieties contain 5 or 6 carbon atoms, examples are *n*-pentyl and *n*-hexyl.

Alkenyl and alkynyl moieties also suitable contain from 2 to 4 carbon atoms in the form of straight or branched chains. Examples are allyl, ethynyl and propargyl.

10 Halo includes fluoro, chloro, bromo and iodo. Most commonly it is fluoro, chloro or bromo and usually fluoro or chloro.

Of particular interest are the compounds of the general formula (1) where X is N and Y is CH (quinolines) and those compounds where X and Y are both N (quinazolin-
lines). Also of interest are those compounds where Y is N and X is CH (isoquinolines).

15 Typically R is halo, for example, chloro.

Typically, R₁ is methyl, ethyl, *n*-propyl, 2,2,2-trifluoromethyl, cyanomethyl, acetylmethyl, methoxycarbonylmethyl, methoxycarbonyl-ethyl, hydroxymethyl, hydroxy-
ethyl, methoxymethyl, methylthiomethyl, ethoxymethyl, 2-methoxyethyl, 2-methylthio-
ethyl, methoxy, ethoxy, *n*-propoxy or *n*-butoxy. Ethyl is a preferred value of R₁ but also
20 of particular interest are methoxy, ethoxy and methoxymethyl.

Typically R₂ is H and at least one, but preferably both of R₃ and R₄ are methyl. When one of R₃ and R₄ is H, the other may be methyl, ethyl or *n*- or *iso*-propyl. When
one of R₃ and R₄ is methyl, the other may be H or ethyl but is preferably also methyl. R₂
also includes C₁₋₄ alkoxymethyl and benzyloxymethyl in which the phenyl ring of the
25 benzyl group optionally carries an alkoxy substituent, e.g. a methoxy substituent. Such
values of R₂ provide compounds of formula (1) that are believed to be pro-pesticidal
compounds:

Typically R₅ is H or methyl, preferably methyl. However, also of particular
interest are compounds where R₅ is hydroxymethyl, methoxymethyl, 1-methoxyethyl and
30 *tert*-butyldimethylsiloxymethyl.

Thus in one particular aspect, the invention provides a compound of the general
formula (1) wherein one of X and Y is N and the other is CH or both of X and Y are N;

R₁ is methyl, ethyl, *n*-propyl, 2,2,2-trifluoromethyl, cyanomethyl, acetylmethyl, methoxycarbonylmethyl, methoxycarbonylethyl, hydroxymethyl, hydroxyethyl, methoxymethyl, methylthiomethyl, ethoxymethyl, 2-methoxyethyl, methoxy, ethoxy, *n*-propoxy or *n*-butoxy; R₂ is H; R₃ and R₄ are both methyl; and R₅ is H, methyl, hydroxymethyl, methoxymethyl or 1-methoxyethyl. Preferably R₁ is ethyl, methoxy, ethoxy or methoxymethyl, especially ethyl. Preferably R₅ is methyl or methoxymethyl.

Compounds that form part of the invention are illustrated in Tables 1 to 57 below.

The compounds in Table 1 are of the general formula (1) where X is N, Y is CH, R₁ is ethyl, R₂ is H, R₃ and R₄ are both methyl and R₅ has the values given in the table.

10 Table 1

Compound No.	R ₅
1	H
2	CH ₃
3	C ₂ H ₅
4	<i>n</i> -C ₃ H ₇
5	<i>i</i> -C ₃ H ₇
6	<i>n</i> -C ₄ H ₉
7	<i>sec</i> -C ₄ H ₉
8	<i>iso</i> -C ₄ H ₉
9	<i>tert</i> -C ₄ H ₉
10	HOCH ₂
11	HOC ₂ H ₄
12	CH ₃ OCH ₂
13	CH ₃ OCH ₂ CH ₂
14	C ₂ H ₅ OCH ₂
15	CH ₃ (CH ₃ O)CH
16	<i>n</i> -C ₃ H ₇ OCH ₂
17	<i>n</i> -C ₃ H ₇ OC ₂ H ₄
18	<i>t</i> -C ₄ H ₉ OCH ₂
19	<i>t</i> -C ₄ H ₉ OC ₂ H ₄
20	CH ₃ SCH ₂

21	$\text{CH}_3\text{SC}_2\text{H}_4$
22	$\text{C}_2\text{H}_5\text{SCH}_2$
23	$\text{C}_2\text{H}_5\text{SC}_2\text{H}_4$
24	$n\text{-C}_3\text{H}_7\text{SCH}_2$
25	$n\text{-C}_4\text{H}_9\text{SCH}_2$
26	$\text{C}_6\text{H}_5\text{OCH}_2$
27	$\text{C}_6\text{H}_5\text{OC}_2\text{H}_4$
28	$4\text{-}t\text{-C}_4\text{H}_9\text{-C}_6\text{H}_4\text{OCH}_2$
29	$4\text{-F-C}_6\text{H}_4\text{OCH}_2$
30	$4\text{-Cl-C}_6\text{H}_4\text{OCH}_2$
31	$4\text{-CH}_3\text{-C}_6\text{H}_4\text{OCH}_2$
32	$4\text{-Br-C}_6\text{H}_4\text{OCH}_2$
33	$2\text{-F-C}_6\text{H}_4\text{OCH}_2$
34	$3,4\text{-Cl}_2\text{-C}_6\text{H}_3\text{OCH}_2$
35	$3\text{-CF}_3\text{-C}_6\text{H}_4\text{OCH}_2$
36	$3,5\text{-Cl}_2\text{-C}_6\text{H}_3\text{OCH}_2$
37	$4\text{-CF}_3\text{O-C}_6\text{H}_5\text{OCH}_2$
38	$2\text{-CF}_3\text{-C}_6\text{H}_4\text{OCH}_2$
39	$4\text{-CF}_3\text{-C}_6\text{H}_4\text{OCH}_2$
40	$2\text{-Br-C}_6\text{H}_4\text{OCH}_2$
41	$2\text{-Cl-C}_6\text{H}_4\text{OCH}_2$
42	$2\text{-CH}_3\text{-4-Cl-C}_6\text{H}_3\text{OCH}_2$
43	$2\text{-CH}_3\text{-5-F-C}_6\text{H}_3\text{OCH}_2$
44	$3\text{-Cl-C}_6\text{H}_4\text{OCH}_2$
45	thien-2-yl-OCH ₂
46	thien-3-yl-OCH ₂
47	$\text{C}_6\text{H}_5\text{CH}_2\text{OCH}_2$
48	thien-2-yl-CH ₂ OCH ₂
49	thien-3-yl-CH ₂ OCH ₂
50	$\text{tert-C}_4\text{H}_9(\text{CH}_3)_2\text{SiOCH}_2$
51	$\text{tert-C}_4\text{H}_9(\text{CH}_3)_2\text{SiOC}_2\text{H}_4$

52	C_6H_5
53	$4\text{-}t\text{-C}_4\text{H}_9\text{-C}_6\text{H}_4$
54	$4\text{-F-C}_6\text{H}_4$
55	$4\text{-Cl-C}_6\text{H}_4$
56	$4\text{-CH}_3\text{-C}_6\text{H}_4$
57	$4\text{-Br-C}_6\text{H}_4$
58	$3\text{CH}_3\text{CO-C}_6\text{H}_4$
59	$3,4\text{-Cl}_2\text{-C}_6\text{H}_3$
60	$3\text{-CF}_3\text{-C}_6\text{H}_4$
61	$3,5\text{-Cl}_2\text{-C}_6\text{H}_3$
62	$4\text{-CF}_3\text{O-C}_6\text{H}_4$
63	$2\text{-CF}_3\text{-C}_6\text{H}_4$
64	$4\text{-CF}_3\text{-C}_6\text{H}_4$
65	$2\text{-Br-C}_6\text{H}_4$
66	$2\text{-Cl-C}_6\text{H}_4$
67	$2\text{-CH}_3\text{-4-Cl-C}_6\text{H}_3$
68	$2\text{-CH}_3\text{-5-F-C}_6\text{H}_3$
69	$3\text{-Cl-C}_6\text{H}_4$
70	thien-2-yl
71	thien-3-yl
72	$\text{C}_6\text{H}_5\text{CH}_2$
73	$4\text{-}t\text{-C}_4\text{H}_9\text{-C}_6\text{H}_4\text{CH}_2$
74	$4\text{-F-C}_6\text{H}_4\text{CH}_2$
75	$4\text{-Cl-C}_6\text{H}_4\text{CH}_2$
76	$4\text{-CH}_3\text{-C}_6\text{H}_4\text{CH}_2$
77	$4\text{-Br-C}_6\text{H}_4\text{CH}_2$
78	$2\text{-F-C}_6\text{H}_4\text{CH}_2$
79	$3,4\text{-Cl}_2\text{-C}_6\text{H}_3\text{CH}_2$
80	$3\text{-CF}_3\text{-C}_6\text{H}_4\text{CH}_2$
81	$3,5\text{-Cl}_2\text{-C}_6\text{H}_3\text{CH}_2$
82	$4\text{-CF}_3\text{O-C}_6\text{H}_5\text{CH}_2$

83	2-CF ₃ -C ₆ H ₄ CH ₂
84	4-CF ₃ -C ₆ H ₄ CH ₂
85	2-Br-C ₆ H ₄ CH ₂
86	2-Cl-C ₆ H ₄ CH ₂
87	2-CH ₃ -4-Cl-C ₆ H ₃ CH ₂
88	2-CH ₃ -5-F-C ₆ H ₃ CH ₂
89	3-Cl-C ₆ H ₄ CH ₂

Table 2

Table 2 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is methyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 2 is the same as compound 1 of Table 1 except that in compound 1 of Table 2 R₁ is methyl instead of ethyl. Similarly, compounds 2 to 89 of Table 2 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 2 R₁ is methyl instead of ethyl.

Table 3

Table 3 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is *n*-propyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 3 is the same as compound 1 of Table 1 except that in compound 1 of Table 3 R₁ is *n*-propyl instead of ethyl. Similarly, compounds 2 to 89 of Table 3 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 3 R₁ is *n*-propyl instead of ethyl.

Table 4

Table 4 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is 2,2,2-trifluoroethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 4 is the same as compound 1 of Table 1 except that in compound 1 of Table 4 R₁ is 2,2,2-trifluoroethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 4 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 4 R₁ is 2,2,2-trifluoroethyl instead of ethyl.

Table 5

Table 5 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is cyanomethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 5 is the same as compound 1 of Table 1 except that in compound 1 of Table 5 R₁ is cyanomethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 5 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 5 R₁ is cyanomethyl instead of ethyl.

Table 6

Table 6 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is acetylmethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 6 is the same as compound 1 of Table 1 except that in compound 1 of Table 6 R₁ is acetylmethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 6 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 2 R₁ is acetylmethyl instead of ethyl.

Table 7

Table 7 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is methoxycarbonylmethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 7 is the same as compound 1 of Table 1 except that in compound 1 of Table 7 R₁ is methoxycarbonylmethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 7 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 7 R₁ is methoxycarbonylmethyl instead of ethyl.

Table 8

Table 8 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is methoxycarbonylethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 8 is the same as compound 1 of Table 1 except that in compound 1 of Table 8 R₁ is methoxycarbonylethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 8 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 8 R₁ is methoxycarbonylethyl instead of ethyl.

Table 9

Table 9 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is hydroxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 9 is the same as compound 1 of Table 1 except
5 that in compound 1 of Table 9 R₁ is hydroxymethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 9 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 9 R₁ is hydroxymethyl instead of ethyl.

Table 10

10 Table 10 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is hydroxyethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 10 is the same as compound 1 of Table 1 except that in compound 1 of Table 10 R₁ is hydroxyethyl instead of ethyl. Similarly, compounds 2
to 89 of Table 10 are the same as compounds 2 to 89 of Table 1, respectively, except that
15 in the compounds of Table 10 R₁ is hydroxyethyl instead of ethyl.

Table 11

Table 11 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is methoxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed
in Table 1. Thus compound 1 of Table 11 is the same as compound 1 of Table 1 except
20 that in compound 1 of Table 11 R₁ is methoxymethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 11 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 11 R₁ is methoxymethyl instead of ethyl.

Table 12

25 Table 12 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is methylthiomethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 12 is the same as compound 1 of Table 1 except that in compound 1 of Table 12 R₁ is methylthiomethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 12 are the same as compounds 2 to 89 of Table 1,
30 respectively, except that in the compounds of Table 12 R₁ is methylthiomethyl instead of ethyl.

Table 13

Table 13 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is ethoxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 13 is the same as compound 1 of Table 1 except that in compound 1 of Table 13 R₁ is ethoxymethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 13 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 13 R₁ is ethoxymethyl instead of ethyl.

Table 14

Table 14 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is 2-methoxyethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 14 is the same as compound 1 of Table 1 except that in compound 1 of Table 14 R₁ is 2-methoxyethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 14 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 14 R₁ is 2-methoxyethyl instead of ethyl.

Table 15

Table 15 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is 2-methylthioethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 15 is the same as compound 1 of Table 1 except that in compound 1 of Table 15 R₁ is 2-methylthioethyl instead of ethyl. Similarly, compounds 2 to 89 of Table 15 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 15 R₁ is 2-methylthioethyl instead of ethyl.

Table 16

Table 16 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is methoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 16 is the same as compound 1 of Table 1 except that in compound 1 of Table 16 R₁ is methoxy instead of ethyl. Similarly, compounds 2 to 89 of Table 16 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 16 R₁ is methoxy instead of ethyl.

Table 17

Table 17 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is ethoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 17 is the same as compound 1 of Table 1 except that in
5 compound 1 of Table 17 R₁ is ethoxy instead of ethyl. Similarly, compounds 2 to 89 of Table 17 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 17 R₁ is ethoxy instead of ethyl.

Table 18

Table 18 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁
10 is *n*-propoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 18 is the same as compound 1 of Table 1 except that in compound 1 of Table 18 R₁ is *n*-propoxy instead of ethyl. Similarly, compounds 2 to 89 of Table 18 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 18 R₁ is *n*-propoxy instead of ethyl.

Table 19

Table 19 consists of 89 compounds of the general formula (1), where X is N, Y is CH, R₁ is *n*-butoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 19 is the same as compound 1 of Table 1 except that
15 in compound 1 of Table 19 R₁ is *n*-butoxy instead of ethyl. Similarly, compounds 2 to 89 of Table 19 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 19 R₁ is *n*-butoxy instead of ethyl.

Table 20

Table 20 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is ethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in
25 Table 1. Thus compound 1 of Table 20 is the same as compound 1 of Table 1 except that in compound 1 of Table 20 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 20 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 20 Y is N instead of CH.

Table 21

Table 21 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is methyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in
30 Table 1. Thus compound 1 of Table 21 is the same as compound 1 of Table 2 except that

in compound 1 of Table 21 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 21 are the same as compounds 2 to 89 of Table 2, respectively, except that in the compounds of Table 21 Y is N instead of CH.

Table 22

- 5 Table 22 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is *n*-propyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 22 is the same as compound 1 of Table 3 except that in compound 1 of Table 22 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 22 are the same as compounds 2 to 89 of Table 3, respectively, except that in the
10 compounds of Table 22 Y is N instead of CH.

Table 23

- Table 23 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is 2,2,2-trifluoroethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 23 is the same as compound 1 of
15 Table 4 except that in compound 1 of Table 23 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 23 are the same as compounds 2 to 89 of Table 4, respectively, except that in the compounds of Table 23 Y is N instead of CH.

Table 24

- Table 24 consists of 89 compounds of the general formula (1), where X and Y are both
20 N, R₁ is cyanomethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 24 is the same as compound 1 of Table 5 except that in compound 1 of Table 24 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 24 are the same as compounds 2 to 89 of Table 5, respectively, except that in the compounds of Table 24 Y is N instead of CH.

Table 25

- Table 25 consists of 89 compounds of the general formula (1), where X and Y are both
25 N, R₁ is acetylmethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 25 is the same as compound 1 of Table 6 except that in compound 1 of Table 25 Y is N instead of CH. Similarly, compounds 2 to
30 89 of Table 25 are the same as compounds 2 to 89 of Table 6, respectively, except that in the compounds of Table 25 Y is N instead of CH.

Table 26

Table 26 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is methoxycarbonylmethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 26 is the same as compound 1 of Table 7 except that in compound 1 of Table 26 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 26 are the same as compounds 2 to 89 of Table 7, respectively, except that in the compounds of Table 26 Y is N instead of CH.

Table 27

Table 27 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is methoxycarbonylethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 27 is the same as compound 1 of Table 8 except that in compound 1 of Table 27 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 27 are the same as compounds 2 to 89 of Table 8, respectively, except that in the compounds of Table 27 Y is N instead of CH.

Table 28

Table 28 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is hydroxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 28 is the same as compound 1 of Table 9 except that in compound 1 of Table 28 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 28 are the same as compounds 2 to 89 of Table 9, respectively, except that in the compounds of Table 28 Y is N instead of CH.

Table 29

Table 29 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is hydroxethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 29 is the same as compound 1 of Table 10 except that in compound 1 of Table 29 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 29 are the same as compounds 2 to 89 of Table 10, respectively, except that in the compounds of Table 29 Y is N instead of CH.

Table 30

Table 30 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is methoxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 30 is the same as compound 1 of Table 11 except that in compound 1 of Table 30 Y is N instead of CH. Similarly, compounds 2 to

89 of Table 30 are the same as compounds 2 to 89 of Table 11, respectively, except that in the compounds of Table 30 Y is N instead of CH.

Table 31

Table 31 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is methylthiomethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 31 is the same as compound 1 of Table 12 except that in compound 1 of Table 31 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 31 are the same as compounds 2 to 89 of Table 12, respectively, except that in the compounds of Table 31 Y is N instead of CH.

Table 32

Table 32 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is ethoxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 32 is the same as compound 1 of Table 13 except that in compound 1 of Table 32 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 32 are the same as compounds 2 to 89 of Table 13, respectively, except that in the compounds of Table 32 Y is N instead of CH.

Table 33

Table 33 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is 2-methoxyethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 33 is the same as compound 1 of Table 14 except that in compound 1 of Table 33 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 33 are the same as compounds 2 to 89 of Table 14, respectively, except that in the compounds of Table 33 Y is N instead of CH.

Table 34

Table 34 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is 2-methylthioethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 34 is the same as compound 1 of Table 15 except that in compound 1 of Table 34 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 34 are the same as compounds 2 to 89 of Table 15, respectively, except that in the compounds of Table 34 Y is N instead of CH.

Table 35

Table 35 consists of 89 compounds of the general formula (1), where X and Y are both N, R₁ is methoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 35 is the same as compound 1 of Table 16 except
5 that in compound 1 of Table 35 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 35 are the same as compounds 2 to 89 of Table 16, respectively, except that in the compounds of Table 35 Y is N instead of CH.

Table 36

Table 36 consists of 89 compounds of the general formula (1), where X and Y are both
10 N, R₁ is ethoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 36 is the same as compound 1 of Table 17 except that in compound 1 of Table 36 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 36 are the same as compounds 2 to 89 of Table 17, respectively, except that in the compounds of Table 36 Y is N instead of CH.

Table 37

Table 37 consists of 89 compounds of the general formula (1), where X and Y are both
15 N, R₁ is *n*-propoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 37 is the same as compound 1 of Table 18 except that in compound 1 of Table 37 Y is N instead of CH. Similarly, compounds 2 to
20 89 of Table 37 are the same as compounds 2 to 89 of Table 18, respectively, except that in the compounds of Table 37 Y is N instead of CH.

Table 38

Table 38 consists of 89 compounds of the general formula (1), where X and Y are both
25 N, R₁ is *n*-butoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 38 is the same as compound 1 of Table 19 except that in compound 1 of Table 38 Y is N instead of CH. Similarly, compounds 2 to 89 of Table 38 are the same as compounds 2 to 89 of Table 19, respectively, except that in the compounds of Table 38 Y is N instead of CH.

Table 39

30 Table 39 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is ethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 39 is the same as compound 1 of Table 1 except that in

compound 1 of Table 39 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 39 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 39 Y is N instead of CH and X is CH instead of N.

5 Table 40

Table 40 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is methyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 40 is the same as compound 1 of Table 2 except that in compound 1 of Table 40 Y is N instead of CH and X is CH instead of N. Similarly,
10 compounds 2 to 89 of Table 40 are the same as compounds 2 to 89 of Table 2, respectively, except that in the compounds of Table 40 Y is N instead of CH and X is CH instead of N.

Table 41

Table 41 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁
15 is *n*-propyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 41 is the same as compound 1 of Table 3 except that in compound 1 of Table 41 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 41 are the same as compounds 2 to 89 of Table 3, respectively, except that in the compounds of Table 41 Y is N instead of CH and X is CH
20 instead of N.

Table 42

Table 42 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is 2,2,2-trifluoroethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 42 is the same as compound 1 of Table 4
25 except that in compound 1 of Table 42 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 42 are the same as compounds 2 to 89 of Table 4, respectively, except that in the compounds of Table 42 Y is N instead of CH and X is CH instead of N.

Table 43

30 Table 43 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is cyanomethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 43 is the same as compound 1 of Table 5 except that

in compound 1 of Table 43 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 43 are the same as compounds 2 to 89 of Table 5, respectively, except that in the compounds of Table 43 Y is N instead of CH and X is CH instead of N.

5 Table 44

Table 44 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is acetylmethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 44 is the same as compound 1 of Table 6 except that in compound 1 of Table 44 Y is N instead of CH and X is CH instead of N. Similarly,
10 compounds 2 to 89 of Table 44 are the same as compounds 2 to 89 of Table 6, respectively, except that in the compounds of Table 44 Y is N instead of CH and X is CH instead of N.

Table 45

Table 45 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁
15 is methoxycarbonylmethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 45 is the same as compound 1 of Table 7 except that in compound 1 of Table 45 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 45 are the same as compounds 2 to 89 of Table 7, respectively, except that in the compounds of Table 45 Y is N instead of CH and
20 X is CH instead of N.

Table 46

Table 46 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is methoxycarbonyl ethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 46 is the same as compound 1 of
25 Table 8 except that in compound 1 of Table 46 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 46 are the same as compounds 2 to 89 of Table 8, respectively, except that in the compounds of Table 46 Y is N instead of CH and X is CH instead of N.

Table 47

30 Table 47 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is hydroxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 47 is the same as compound 1 of Table 9 except

that in compound 1 of Table 47 Y is N instead of CH and X is CH instead of N.

Similarly, compounds 2 to 89 of Table 47 are the same as compounds 2 to 89 of Table 9, respectively, except that in the compounds of Table 47 Y is N instead of CH and X is CH instead of N.

5 Table 48

Table 48 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is hydroxethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 48 is the same as compound 1 of Table 10 except that in compound 1 of Table 48 Y is N instead of CH and X is CH instead of N.

10 Similarly, compounds 2 to 89 of Table 48 are the same as compounds 2 to 89 of Table 10, respectively, except that in the compounds of Table 48 Y is N instead of CH and X is CH instead of N.

Table 49

15 Table 49 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is methoxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 49 is the same as compound 1 of Table 11 except that in compound 1 of Table 49 Y is N instead of CH and X is CH instead of N.

 Similarly, compounds 2 to 89 of Table 49 are the same as compounds 2 to 89 of Table 11, respectively, except that in the compounds of Table 49 Y is N instead of CH and X is
20 CH instead of N.

Table 50

 Table 50 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is methylthiomethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 50 is the same as compound 1 of Table 12
25 except that in compound 1 of Table 50 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 50 are the same as compounds 2 to 89 of Table 12, respectively, except that in the compounds of Table 50 Y is N instead of CH and X is CH instead of N.

Table 51

30 Table 51 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is ethoxymethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 51 is the same as compound 1 of Table 13 except

that in compound 1 of Table 51 Y is N instead of CH and X is CH instead of N.

Similarly, compounds 2 to 89 of Table 51 are the same as compounds 2 to 89 of Table 13, respectively, except that in the compounds of Table 51 Y is N instead of CH and X is CH instead of N.

5 Table 52

Table 52 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is 2-methoxyethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 52 is the same as compound 1 of Table 14 except that in compound 1 of Table 52 Y is N instead of CH and X is CH instead of N.

10 Similarly, compounds 2 to 89 of Table 52 are the same as compounds 2 to 89 of Table 14, respectively, except that in the compounds of Table 52 Y is N instead of CH and X is CH instead of N.

Table 53

Table 53 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is 2-methylthioethyl, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 53 is the same as compound 1 of Table 15 except that in compound 1 of Table 53 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 53 are the same as compounds 2 to 89 of Table 15, respectively, except that in the compounds of Table 53 Y is N instead of CH and X is CH instead of N.

Table 54

Table 54 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is methoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 54 is the same as compound 1 of Table 16 except that in compound 1 of Table 54 Y is N instead of CH and X is CH instead of N.

Similarly, compounds 2 to 89 of Table 54 are the same as compounds 2 to 89 of Table 1, respectively, except that in the compounds of Table 16 Y is N instead of CH and X is CH instead of N.

Table 55

30 Table 55 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is ethoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 55 is the same as compound 1 of Table 17 except that in

compound 1 of Table 55 Y is N instead of CH and X is CH instead of N. Similarly, compounds 2 to 89 of Table 55 are the same as compounds 2 to 89 of Table 17, respectively, except that in the compounds of Table 55 Y is N instead of CH and X is CH instead of N.

5 Table 56

Table 56 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is *n*-propoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 56 is the same as compound 1 of Table 18 except that in compound 1 of Table 56 Y is N instead of CH and X is CH instead of N.

10 Similarly, compounds 2 to 89 of Table 56 are the same as compounds 2 to 89 of Table 18, respectively, except that in the compounds of Table 56 Y is N instead of CH and X is CH instead of N.

Table 57

Table 57 consists of 89 compounds of the general formula (1), where X is CH, Y is N, R₁ is *n*-butoxy, R₂ is hydrogen, R₃ and R₄ are both methyl and R₅ has the values listed in Table 1. Thus compound 1 of Table 57 is the same as compound 1 of Table 19 except that in compound 1 of Table 57 Y is N instead of CH and X is CH instead of N.


Similarly, compounds 2 to 89 of Table 57 are the same as compounds 2 to 89 of Table 19, respectively, except that in the compounds of Table 57 Y is N instead of CH and X is CH instead of N.

20 The compounds of formula (1) may be prepared as outlined in Schemes 1 to 9 below in which X, Y, R₁, R₂, R₃, R₄ and R₅ have the meanings given above, R₆ is straight-chain C₁₋₄ alkyl, R₇ and R₈ are independently H or C₁₋₄ alkyl, L is a leaving group such as a halide, for example iodide, an alkyl or aryl sulphonyloxy group, for example methylsulphonyloxy and tosyloxy or a triflate, Hal is halogen, R_a is hydrogen or C₁₋₃ alkyl, R_b is hydrogen or C₁₋₃ alkyl, provided that the total number of carbon atoms in R_a and R_b do not exceed three, R_c is C₁₋₆ alkyl, optionally substituted benzyl or optionally substituted thienylmethyl and R_d has the meaning ascribed to it in the text.

As shown in Scheme 1, the compounds of general formula (1) may be prepared by reacting a compound of the general formula (2) with a compound of the general formula (3) in the presence of a base in a suitable solvent. Typical solvents include *N,N*-dimethyl-


formamide and *N*-methylpyrrolidin-2-one. Suitable bases include potassium carbonate, sodium hydride or *diisopropylethylamine*.

Scheme 1



As shown in Scheme 2, compounds of the general formula (3) may be prepared by reacting an amine of the general formula (5) with an acid halide of the general formula (4), or the corresponding acid anhydride, in the presence of a suitable inorganic or organic base, such as potassium carbonate or *diisopropylethylamine*, in a solvent such as dichloromethane or tetrahydrofuran.

Scheme 2



10


As shown in Scheme 3, amines of the general formula (5), wherein R₂ is H, correspond to amines of the general formula (9) and may be prepared by alkylation of a silyl-protected aminoalkyne of the general formula (7) using a suitable base, such as *n*-butyl lithium, followed by reaction with a suitable alkylating reagent R₅L, such as an alkyl iodide, for example, methyl iodide, to form an alkylated compound of the general formula (8). In a similar procedure, a silyl-protected aminoalkyne of the general formula (7) may be reacted with a carbonyl derivative R_aCOR_b, for example formaldehyde, using a suitable base, such as *n*-butyl lithium, to provide an aminoalkyne (8) containing a hydroxyalkyl moiety. The silyl protecting group may then be removed from a compound of the general formula (8) with, for example, an aqueous acid to form an aminoalkyne of the general formula (9). Aminoalkynes of the general formula (9) may be further derivatised, for instance when R⁵ is a hydroxyalkyl group, for example, by reacting a compound of the general formula (9) with a silylating agent, for example *t*-butyldimethyl-

15

20

silyl chloride, to give a derivative silylated on oxygen of the general formula (9a). In addition, a compound of the general formula (9) may be treated with a base, such as sodium hydride or potassium *bis*(trimethylsilyl)amide followed by a compound R_cL to give a compound of the general formula (9b). In an alternative sequence, a compound of general formula (8) may be treated with a base, such as sodium or potassium *bis*(trimethylsilyl)amide, followed by a compound R_cL , where L represents a halogen or sulphonate ester such as OSO_2Me , or OSO_2-4 -tolyl, for example ethyl iodide, to give, after removal of the silyl protecting group, compounds of general formula (9b).

10 Scheme 3



Silyl-protected aminoalkynes of the general formula (7) may be obtained by reacting amines of general formula (6) with 1,2-*bis*-(chlorodimethylsilyl)ethane in the


presence of a suitable base, such as a tertiary organic amine base, for example, triethylamine.

Amines of the general formula (6) are either commercially available or may be prepared by standard literature methods (see, for example, EP-A-0834498).

- 5 : Alternatively, as shown in Scheme 4, compounds of the general formula (1) may be prepared by condensing a compound of the general formula (11), wherein R is H with an amine of the general formula (5) using suitable activating reagents such as 1-hydroxybenzotriazole and *N*-(3-dimethylaminopropyl)-*N'*-ethyl-carbodiimide hydrochloride.

- 10 Where R₂ is other than hydrogen, the R₂ group may be introduced into an aminoalkyne of the general formula (9) by known techniques to form an amine of the general formula (5).

Scheme 4




Compounds of the general formula (12) may be prepared by the hydrolysis of the corresponding esters of general formula (11), wherein R_d is C_{1-4} alkyl, using known techniques. The esters of the general formula (11), wherein R_d is C_{1-4} alkyl and also acids of the general formula (11), wherein R_d is H, may be prepared by reacting a compound of
5 the general formula (2) with an ester or acid of the general formula (10a) in the presence of a suitable base, such as potassium carbonate or sodium hydride, in a suitable solvent, such as *N,N*-dimethylformamide. The esters or acids of the general formula (10a) are either commercially available or may be prepared by standard literature methods from commercially available materials.

10 Alternatively, as shown in Scheme 4, compounds of the general formula (11) may be prepared under Mitsunobu conditions by reacting a compound of the general formula (2) with a compound of the general formula (10b), wherein R_d is C_{1-4} alkyl, using a phosphine, such as triphenyl phosphine, and an azoester, such as diethyl azodicarboxylate.


15 Similarly, compounds of the general formula (1) may be prepared by reacting a compound of general formula (10d) with a compound of the general formula (2) under Mitsunobu conditions using a phosphine, such as triphenyl phosphine, and an azoester, such as diethyl azodicarboxylate. Compounds of general formula (10d) may be prepared from a compound of general formula (10c) and an amine of general formula (5) using
20 suitable activating reagents such as 1-hydroxybenzotriazole and *N*-(3-dimethylamino-propyl)-*N'*-ethyl-carbodiimide hydrochloride. Compounds (10b) and (10c) are either known compounds or may be made from known compounds.

In another method, the compounds of the general formula (1) may be prepared by reacting an acid halide of the general formula (13) with an amine of the general formula
25 (5) in a suitable solvent, such as dichloromethane, in the presence of a tertiary amine, such as triethylamine, and an activating agent, such as 4-dimethylaminopyridine.

As shown in Scheme 5, an acid halide of the general formula (13) may be prepared by chlorinating a compound of the general formula (12) with a suitable chlorinating agent, such as oxalyl chloride, in a suitable solvent, such as dichloro-
30 methane, and in the presence of, for example, *N,N*-dimethylformamide. The compounds of the general formula (12) correspond to the compounds of general formula (11), wherein R is H.

Scheme 5

5 As shown in Scheme 6, compounds of the general formula (1), wherein R_5 is H, may be reacted under Sonogashira conditions with, for example, optionally substituted phenyl or thienyl chlorides, bromides, iodides or triflates to form substituted phenyl or thienyl compounds of general formula (1), wherein R_5 is an optionally substituted phenyl or thienyl group. A suitable palladium catalyst is tetrakis(triphenylphosphine)-
10 palladium(0).


Scheme 6

Compounds of the general formula (1) wherein R_1 is straight-chain C_{1-4} alkoxy, such as compounds of the general formula (14) wherein R_6 is as defined above, may be prepared as shown in Scheme 7. Thus, esters of the formula (15) may be halogenated to give haloesters of the general formula (16), by treatment with a suitable halogenating agent, such as N-bromosuccinimide, in a suitable solvent such as carbon tetrachloride, at between ambient temperature and the reflux temperature of the solvent. The haloesters of
15 the general formula (16) can be reacted with an alkali metal compound M^+OR_6 , where M is suitably sodium or potassium in, for example, an alcohol R_6OH as solvent, at between $0^\circ C$ and $40^\circ C$, preferably at ambient temperature, to give compounds of the general formula (17). The esters (17) can be hydrolysed to acids of the general formula (18), by
20

treatment with an alkali metal hydroxide, such as sodium hydroxide, in an aqueous alcohol R_6OH , at between ambient temperature and reflux.


A carboxylic acid of the general formula (18) can be condensed with an amine of the general formula (5) to give a compound of the general formula (14), where R_6 is as defined above, using suitable activating reagents such as 1-hydroxybenzotriazole and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride.

Scheme 7



Compounds of the general formula (1), wherein R_1 is C_{1-4} alkyl, C_{3-4} alkenyl, C_{3-4} alkynyl, or an alkoxyalkyl group where the total number of carbon atoms is 2 or 3, may be prepared as shown in Scheme 8. Thus, the substituted acetic acid (19) may be treated with at least two equivalents of a base, such as lithium diisopropylamide, in a suitable solvent such as tetrahydrofuran, at a temperature between $-78^\circ C$ and ambient temperature, with an alkylating agent such as R_1L to give carboxylic acids of the general formula (20) upon acidification.


Scheme 8



As shown in Scheme 9, compounds of the general formula (1), where R_1 is a C_{3-6} alkenyl group, may be prepared from esters of the general formula (21), wherein R_7 and

R_8 are as defined above. Esters of the general formula (21) are treated with a strong base, such as lithium *bis*(trimethylsilyl)amide, at between -78°C and ambient temperature, preferably at -78°C , and then reacted with a trialkylsilyl chloride $R_3\text{SiCl}$, such as trimethylsilyl chloride, or trialkylsilyl triflate $R_3\text{SiOSO}_2\text{CF}_3$, and allowed to warm to ambient temperature. The resultant acids of the general formula (22) obtained after hydrolysis can be condensed with amines of the general formula (5) to give the compounds of the general formula (23), using suitable activating reagents such as 1-hydroxybenzotriazole and *N*-(3-dimethylaminopropyl)-*N'*-ethyl-carbodiimide hydrochloride.

10 Scheme 9



Other compounds of the invention may be prepared by transforming the substituents in the compounds of the general formula (1) using known procedures e.g. by the alkylation of compounds of the general formula (1), wherein R_2 is H or R_5 is H.

The compounds of formula (1) are active fungicides and may be used to control one or more of the following pathogens: *Pyricularia oryzae* (*Magnaporthe grisea*) on rice and wheat and other *Pyricularia* spp. on other hosts; *Puccinia triticina* (or *recondita*), *Puccinia striiformis* and other rusts on wheat, *Puccinia hordei*, *Puccinia striiformis* and other rusts on barley, and rusts on other hosts (for example turf, rye, coffee, pears, apples, peanuts, sugar beet, vegetables and ornamental plants); *Erysiphe cichoracearum* on cucurbits (for example melon); *Blumeria* (or *Erysiphe*) *graminis* (powdery mildew) on barley, wheat, rye and turf and other powdery mildews on various hosts, such as *Sphaerotheca macularis* on hops, *Sphaerotheca fusca* (*Sphaerotheca fuliginea*) on cucurbits (for example cucumber), *Leveillula taurica* on tomatoes, aubergine and green pepper, *Podosphaera leucotricha* on apples and *Uncinula necator* on vines; *Cochliobolus* spp., *Helminthosporium* spp., *Drechslera* spp. (*Pyrenophora* spp.), *Rhynchosporium* spp., *Mycosphaerella graminicola* (*Septoria tritici*) and *Phaeosphaeria nodorum*.

(*Stagonospora nodorum* or *Septoria nodorum*), *Pseudocercospora herpotrichoides* and *Gaeumannomyces graminis* on cereals (for example wheat, barley, rye), turf and other hosts; *Cercospora arachidicola* and *Cercosporidium personatum* on peanuts and other *Cercospora* spp. on other hosts, for example sugar beet, bananas, soya beans and rice;

5 *Botrytis cinerea* (grey mould) on tomatoes, strawberries, vegetables, vines and other hosts and other *Botrytis* spp. on other hosts; *Alternaria* spp. on vegetables (for example carrots), oil-seed rape, apples, tomatoes, potatoes, cereals (for example wheat) and other hosts; *Venturia* spp. (including *Venturia inaequalis* (scab)) on apples, pears, stone fruit, tree nuts and other hosts; *Cladosporium* spp. on a range of hosts including cereals (for

10 example wheat) and tomatoes; *Monilinia* spp. on stone fruit, tree nuts and other hosts; *Didymella* spp. on tomatoes, turf, wheat, cucurbits and other hosts; *Phoma* spp. on oil-seed rape, turf, rice, potatoes, wheat and other hosts; *Aspergillus* spp. and *Aureobasidium* spp. on wheat, lumber and other hosts; *Ascochyta* spp. on peas, wheat, barley and other hosts; *Stemphylium* spp. (*Pleospora* spp.) on apples, pears, onions and

15 other hosts; summer diseases (for example bitter rot (*Glomerella cingulata*), black rot or frog-eye leaf spot (*Botryosphaeria obtusa*), Brooks fruit spot (*Mycosphaerella pomi*), Cedar apple rust (*Gymnosporangium juniperi-virginianae*), sooty blotch (*Gloeodes pomigena*), flyspeck (*Schizothyrium pomi*) and white rot (*Botryosphaeria dothidea*)) on apples and pears; *Plasmopara viticola* on vines; other downy mildews, such as *Bremia*

20 *lactucae* on lettuce, *Peronospora* spp. on soybeans, tobacco, onions and other hosts, *Pseudoperonospora humuli* on hops and *Pseudoperonospora cubensis* on cucurbits; *Pythium* spp. (including *Pythium ultimum*) on turf and other hosts; *Phytophthora* *infestans* on potatoes and tomatoes and other *Phytophthora* spp. on vegetables, strawberries, avocado, pepper, ornamentals, tobacco, cocoa and other hosts;

25 *Thanatephorus cucumeris* on rice and turf and other *Rhizoctonia* spp. on various hosts such as wheat and barley, peanuts, vegetables, cotton and turf; *Sclerotinia* spp. on turf, peanuts, potatoes, oil-seed rape and other hosts; *Sclerotium* spp. on turf, peanuts and other hosts; *Gibberella fujikuroi* on rice; *Colletotrichum* spp. on a range of hosts including turf, coffee and vegetables; *Laetisaria fuciformis* on turf; *Mycosphaerella* spp. on bananas,

30 peanuts, citrus, pecans, papaya and other hosts; *Diaporthe* spp. on citrus, soybean, melon, pears, lupin and other hosts; *Elsinoe* spp. on citrus, vines, olives, pecans, roses and other hosts; *Verticillium* spp. on a range of hosts including hops, potatoes and tomatoes;

Pyrenopeziza spp. on oil-seed rape and other hosts; *Oncobasidium theobromae* on cocoa causing vascular streak dieback; *Fusarium* spp., *Typhula* spp., *Microdochium nivale*, *Ustilago* spp., *Urocystis* spp., *Tilletia* spp. and *Claviceps purpurea* on a variety of hosts but particularly wheat, barley, turf and maize; *Ramularia* spp. on sugar beet, barley and other hosts; post-harvest diseases particularly of fruit (for example *Penicillium digitatum*, *Penicillium italicum* and *Trichoderma viride* on oranges, *Colletotrichum musae* and *Gloeosporium musarum* on bananas and *Botrytis cinerea* on grapes); other pathogens on vines, notably *Eutypa lata*, *Guignardia bidwellii*, *Phellinus igniarius*, *Phomopsis viticola*, *Pseudopeziza tracheiphila* and *Stereum hirsutum*; other pathogens on trees (for example *Lophodermium seeditiosum*) or lumber, notably *Cephalosporium fragrans*, *Ceratocystis* spp., *Ophiostoma piceae*, *Penicillium* spp., *Trichoderma pseudokoningii*, *Trichoderma viride*, *Trichoderma harzianum*, *Aspergillus niger*, *Leptographium lindbergii* and *Aureobasidium pullulans*; and fungal vectors of viral diseases (for example *Polymyxa graminis* on cereals as the vector of barley yellow mosaic virus (BYMV) and *Polymyxa betae* on sugar beet as the vector of rhizomania).

The compounds of formula (I) show particularly good activity against the Oomycete class of pathogens such as *Phytophthora infestans*, *Plasmopara* species, e.g. *Plasmopara viticola* and *Pythium* species e.g. *Pythium ultimum*.

A compound of formula (1) may move acropetally, basipetally or locally in plant tissue to be active against one or more fungi. Moreover, a compound of formula (1) may be volatile enough to be active in the vapour phase against one or more fungi on the plant.

The invention therefore provides a method of combating or controlling phytopathogenic fungi which comprises applying a fungicidally effective amount of a compound of formula (1), or a composition containing a compound of formula (1), to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or any other plant growth medium, e.g. nutrient solution.

The term "plant" as used herein includes seedlings, bushes and trees. Furthermore, the fungicidal method of the invention includes protectant, curative, systemic, eradicator and antispore treatments.

The compounds of formula (1) are preferably used for agricultural, horticultural and turfgrass purposes in the form of a composition.

In order to apply a compound of formula (1) to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or any other growth medium, a compound of formula (1) is usually formulated into a composition which includes, in addition to the compound of formula (1), a suitable inert diluent or carrier and, optionally, a surface active agent (SFA). SFAs are chemicals which are able to modify the properties of an interface (for example, liquid/solid, liquid/air or liquid/liquid interfaces) by lowering the interfacial tension and thereby leading to changes in other properties (for example dispersion, emulsification and wetting). It is preferred that all compositions (both solid and liquid formulations) comprise, by weight, 0.0001 to 95%, more preferably 1 to 85%, for example 5 to 60%, of a compound of formula (1). The composition is generally used for the control of fungi such that a compound of formula (1) is applied at a rate of from 0.1g to 10kg per hectare, preferably from 1g to 6kg per hectare, more preferably from 1g to 1kg per hectare.

When used in a seed dressing, a compound of formula (1) is used at a rate of 0.0001g to 10g (for example 0.001g or 0.05g), preferably 0.005g to 10g, more preferably 0.005g to 4g, per kilogram of seed.

In another aspect the present invention provides a fungicidal composition comprising a fungicidally effective amount of a compound of formula (1) and a suitable carrier or diluent therefor.

In a still further aspect the invention provides a method of combating and controlling fungi at a locus, which comprises treating the fungi, or the locus of the fungi with a fungicidally effective amount of a composition comprising a compound of formula (1).

The compositions can be chosen from a number of formulation types, including dustable powders (DP), soluble powders (SP), water soluble granules (SG), water dispersible granules (WG), wettable powders (WP), granules (GR) (slow or fast release), soluble concentrates (SL), oil miscible liquids (OL), ultra low volume liquids (UL), emulsifiable concentrates (EC), dispersible concentrates (DC), emulsions (both oil in water (EW) and water in oil (EO)), micro-emulsions (ME), suspension concentrates (SC), aerosols, fogging/smoke formulations, capsule suspensions (CS) and seed treatment formulations. The formulation type chosen in any instance will depend upon the

particular purpose envisaged and the physical, chemical and biological properties of the compound of formula (1).

Dustable powders (DP) may be prepared by mixing a compound of formula (1) with one or more solid diluents (for example natural clays, kaolin, pyrophyllite, bentonite, alumina, montmorillonite, kieselguhr, chalk, diatomaceous earths, calcium phosphates, calcium and magnesium carbonates, sulphur, lime, flours, talc and other organic and inorganic solid carriers) and mechanically grinding the mixture to a fine powder.

Soluble powders (SP) may be prepared by mixing a compound of formula (1) with one or more water-soluble inorganic salts (such as sodium bicarbonate, sodium carbonate or magnesium sulphate) or one or more water-soluble organic solids (such as a polysaccharide) and, optionally, one or more wetting agents, one or more dispersing agents or a mixture of said agents to improve water dispersibility/solubility. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water soluble granules (SG).

Wettable powders (WP) may be prepared by mixing a compound of formula (1) with one or more solid diluents or carriers, one or more wetting agents and, preferably, one or more dispersing agents and, optionally, one or more suspending agents to facilitate the dispersion in liquids. The mixture is then ground to a fine powder. Similar compositions may also be granulated to form water dispersible granules (WG).

Granules (GR) may be formed either by granulating a mixture of a compound of formula (1) and one or more powdered solid diluents or carriers, or from pre-formed blank granules by absorbing a compound of formula (1) (or a solution thereof, in a suitable agent) in a porous granular material (such as pumice, attapulgate clays, fuller's earth, kieselguhr, diatomaceous earths or ground corn cobs) or by adsorbing a compound of formula (1) (or a solution thereof, in a suitable agent) on to a hard core material (such as sands, silicates, mineral carbonates, sulphates or phosphates) and drying if necessary. Agents which are commonly used to aid absorption or adsorption include solvents (such as aliphatic and aromatic petroleum solvents, alcohols, ethers, ketones and esters) and sticking agents (such as polyvinyl acetates, polyvinyl alcohols, dextrans, sugars and vegetable oils). One or more other additives may also be included in granules (for example an emulsifying agent, wetting agent or dispersing agent).

Dispersible Concentrates (DC) may be prepared by dissolving a compound of formula (1) in water or an organic solvent, such as a ketone, alcohol or glycol ether. These solutions may contain a surface active agent (for example to improve water dilution or prevent crystallisation in a spray tank).

5 Emulsifiable concentrates (EC) or oil-in-water emulsions (EW) may be prepared by dissolving a compound of formula (1) in an organic solvent (optionally containing one or more wetting agents, one or more emulsifying agents or a mixture of said agents). Suitable organic solvents for use in ECs include aromatic hydrocarbons (such as alkylbenzenes or alkylnaphthalenes, exemplified by SOLVESSO 100, SOLVESSO 150
10 and SOLVESSO 200; SOLVESSO is a Registered Trade Mark), ketones (such as cyclohexanone or methylcyclohexanone), alcohols (such as benzyl alcohol, furfuryl alcohol or butanol), *N*-alkylpyrrolidones (such as *N*-methylpyrrolidone or *N*-octylpyrrolidone), dimethyl amides of fatty acids (such as C₈-C₁₀ fatty acid dimethylamide) and chlorinated hydrocarbons. An EC product may spontaneously
15 emulsify on addition to water, to produce an emulsion with sufficient stability to allow spray application through appropriate equipment. Preparation of an EW involves obtaining a compound of formula (1) either as a liquid (if it is not a liquid at room temperature, it may be melted at a reasonable temperature, typically below 70°C) or in solution (by dissolving it in an appropriate solvent) and then emulsifying the resultant
20 liquid or solution into water containing one or more SFAs, under high shear, to produce an emulsion. Suitable solvents for use in EWs include vegetable oils, chlorinated hydrocarbons (such as chlorobenzenes), aromatic solvents (such as alkylbenzenes or alkylnaphthalenes) and other appropriate organic solvents which have a low solubility in water.

25 Microemulsions (ME) may be prepared by mixing water with a blend of one or more solvents with one or more SFAs, to produce spontaneously a thermodynamically stable isotropic liquid formulation. A compound of formula (1) is present initially in either the water or the solvent/SFA blend. Suitable solvents for use in MEs include those hereinbefore described for use in in ECs or in EWs. An ME may be either an oil-in-water
30 or a water-in-oil system (which system is present may be determined by conductivity measurements) and may be suitable for mixing water-soluble and oil-soluble pesticides in

the same formulation. An ME is suitable for dilution into water, either remaining as a microemulsion or forming a conventional oil-in-water emulsion.

Suspension concentrates (SC) may comprise aqueous or non-aqueous suspensions of finely divided insoluble solid particles of a compound of formula (1). SCs may be prepared by ball or bead milling the solid compound of formula (1) in a suitable medium, optionally with one or more dispersing agents, to produce a fine particle suspension of the compound. One or more wetting agents may be included in the composition and a suspending agent may be included to reduce the rate at which the particles settle. Alternatively, a compound of formula (1) may be dry milled and added to water, containing agents hereinbefore described, to produce the desired end product.

Aerosol formulations comprise a compound of formula (1) and a suitable propellant (for example *n*-butane). A compound of formula (1) may also be dissolved or dispersed in a suitable medium (for example water or a water miscible liquid, such as *n*-propanol) to provide compositions for use in non-pressurised, hand-actuated spray pumps.

A compound of formula (1) may be mixed in the dry state with a pyrotechnic mixture to form a composition suitable for generating, in an enclosed space, a smoke containing the compound.

Capsule suspensions (CS) may be prepared in a manner similar to the preparation of EW formulations but with an additional polymerisation stage such that an aqueous dispersion of oil droplets is obtained, in which each oil droplet is encapsulated by a polymeric shell and contains a compound of formula (1) and, optionally, a carrier or diluent therefor. The polymeric shell may be produced by either an interfacial polycondensation reaction or by a coacervation procedure. The compositions may provide for controlled release of the compound of formula (1) and they may be used for seed treatment. A compound of formula (1) may also be formulated in a biodegradable polymeric matrix to provide a slow, controlled release of the compound.

A composition may include one or more additives to improve the biological performance of the composition (for example by improving wetting, retention or distribution on surfaces; resistance to rain on treated surfaces; or uptake or mobility of a compound of formula (1)). Such additives include surface active agents, spray additives based on oils, for example certain mineral oils or natural plant oils (such as soy bean and

rape seed oil), and blends of these with other bio-enhancing adjuvants (ingredients which may aid or modify the action of a compound of formula (1)).

A compound of formula (1) may also be formulated for use as a seed treatment, for example as a powder composition, including a powder for dry seed treatment (DS), a water soluble powder (SS) or a water dispersible powder for slurry treatment (WS), or as
5 a liquid composition, including a flowable concentrate (FS), a solution (LS) or a capsule suspension (CS). The preparations of DS, SS, WS, FS and LS compositions are very similar to those of, respectively, DP, SP, WP, SC and DC compositions described above. Compositions for treating seed may include an agent for assisting the adhesion of the
10 composition to the seed (for example a mineral oil or a film-forming barrier).

Wetting agents, dispersing agents and emulsifying agents may be SFAs of the cationic, anionic, amphoteric or non-ionic type.

Suitable SFAs of the cationic type include quaternary ammonium compounds (for example cetyltrimethyl ammonium bromide), imidazolines and amine salts.

15 Suitable anionic SFAs include alkali metals salts of fatty acids, salts of aliphatic monoesters of sulphuric acid (for example sodium lauryl sulphate), salts of sulphonated aromatic compounds (for example sodium dodecylbenzenesulphonate, calcium dodecylbenzenesulphonate, butylnaphthalene sulphonate and mixtures of sodium di-*isopropyl*- and tri-*isopropyl*-naphthalene sulphonates), ether sulphates, alcohol ether sulphates (for
20 example sodium laureth-3-sulphate), ether carboxylates (for example sodium laureth-3-carboxylate), phosphate esters (products from the reaction between one or more fatty alcohols and phosphoric acid (predominately mono-esters) or phosphorus pentoxide (predominately di-esters), for example the reaction between lauryl alcohol and tetraphosphoric acid; additionally these products may be ethoxylated), sulphosuccinamates,
25 paraffin or olefine sulphonates, taurates and lignosulphonates.

Suitable SFAs of the amphoteric type include betaines, propionates and glycines.

Suitable SFAs of the non-ionic type include condensation products of alkylene oxides, such as ethylene oxide, propylene oxide, butylene oxide or mixtures thereof, with
30 fatty alcohols (such as oleyl alcohol or cetyl alcohol) or with alkylphenols (such as octylphenol, nonylphenol or octylcresol); partial esters derived from long chain fatty acids or hexitol anhydrides; condensation products of said partial esters with ethylene oxide;

block polymers (comprising ethylene oxide and propylene oxide); alkanolamides; simple esters (for example fatty acid polyethylene glycol esters); amine oxides (for example lauryl dimethyl amine oxide); and lecithins.

Suitable suspending agents include hydrophilic colloids (such as polysaccharides, polyvinylpyrrolidone or sodium carboxymethylcellulose) and swelling clays (such as bentonite or attapulgate).

A compound of formula (1) may be applied by any of the known means of applying fungicidal compounds. For example, it may be applied, formulated or unformulated, to any part of the plant, including the foliage, stems, branches or roots, to the seed before it is planted or to other media in which plants are growing or are to be planted (such as soil surrounding the roots, the soil generally, paddy water or hydroponic culture systems), directly or it may be sprayed on, dusted on, applied by dipping, applied as a cream or paste formulation, applied as a vapour or applied through distribution or incorporation of a composition (such as a granular composition or a composition packed in a water-soluble bag) in soil or an aqueous environment.

A compound of formula (1) may also be injected into plants or sprayed onto vegetation using electrodynamic spraying techniques or other low volume methods, or applied by land or aerial irrigation systems.

Compositions for use as aqueous preparations (aqueous solutions or dispersions) are generally supplied in the form of a concentrate containing a high proportion of the active ingredient, the concentrate being added to water before use. These concentrates, which may include DCs, SCs, ECs, EWs, MEs SGs, SPs, WPs, WGs and CSs, are often required to withstand storage for prolonged periods and, after such storage, to be capable of addition to water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. Such aqueous preparations may contain varying amounts of a compound of formula (1) (for example 0.0001 to 10%, by weight) depending upon the purpose for which they are to be used.

A compound of formula (1) may be used in mixtures with fertilisers (for example nitrogen-, potassium- or phosphorus-containing fertilisers). Suitable formulation types include granules of fertiliser. The mixtures suitably contain up to 25% by weight of the compound of formula (1).

The invention therefore also provides a fertiliser composition comprising a fertiliser and a compound of formula (1).


The compositions of this invention may contain other compounds having biological activity, for example micronutrients or compounds having similar or
5 complementary fungicidal activity or which possess plant growth regulating, herbicidal, insecticidal, nematocidal or acaricidal activity.

By including another fungicide, the resulting composition may have a broader spectrum of activity or a greater level of intrinsic activity than the compound of formula (1) alone. Further the other fungicide may have a synergistic effect on the fungicidal
10 activity of the compound of formula (1).

The compound of formula (1) may be the sole active ingredient of the composition or it may be admixed with one or more additional active ingredients such as a pesticide, fungicide, synergist, herbicide or plant growth regulator where appropriate. An additional active ingredient may: provide a composition having a broader spectrum of
15 activity or increased persistence at a locus; synergise the activity or complement the activity (for example by increasing the speed of effect or overcoming repellency) of the compound of formula (1); or help to overcome or prevent the development of resistance to individual components. The particular additional active ingredient will depend upon the intended utility of the composition.

20 Examples of fungicidal compounds which may be included in the composition of the invention are AC 382042 (*N*-(1-cyano-1,2-dimethylpropyl)-2-(2,4-dichlorophenoxy) propionamide), acibenzolar-S-methyl, alanycarb, aldimorph, anilazine, azaconazole, azafenidin, azoxystrobin, benalaxyl, benomyl, benthiavalicarb, biloxazol, bitertanol, blasticidin S, boscalid (new name for nicobifen), bromuconazole, bupirimate, captafol,
25 captan, carbendazim, carbendazim chlorhydrate, carboxin, carpropamid, carvone, CGA 41396, CGA 41397, chinomethionate, chlorbenzthiazole, chlorothalonil, chlorozoline, clozylacon, copper containing compounds such as copper oxychloride, copper oxyquino- late, copper sulphate, copper tallate, and Bordeaux mixture, cyamidazosulfamid, cyazofamid (IKF-916), cyflufenamid, cymoxanil, cyproconazole, cyprodinil, debacarb,
30 di-2-pyridyl disulphide 1,1'-dioxide, dichlofluanid, diclocymet, diclomezine, dicloran, diethofencarb, difenoconazole, difenzoquat, diflumetorim, *O,O*-di-*iso*-propyl-*S*-benzyl thiophosphate, dimefluazole, dimetconazole, dimethirimol, dimethomorph,

dimoxystrobin, diniconazole, dinocap, dithianon, dodecyl dimethyl ammonium chloride, dodemorph, dodine, doguadine, edifenphos, epoxiconazole, ethaboxam, ethirimol, ethyl (Z)-N-benzyl-N([methyl(methyl-thioethylideneaminooxycarbonyl)amino]thio)- β -alaninate, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, 5 fenhexamid, fenoxanil (AC 382042), fencpiclonil, fenpropidin, fenpropimorph, fentin acetate, fentin hydroxide, ferbam, ferimzone, fluazinam, fludioxonil, flumetover, flumorph, fluoroimide, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, fosetyl-aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, hexaconazole, hydroxyisoxazole, hymexazole, imazalil, imibenconazole, 10 iminoctadine, iminoctadine triacetate, ipconazole, iprobenfos, iprodione, iprovalicarb, isopropanyl butyl carbamate, isoprothiolane, kasugamycin, kresoxim-methyl, LY186054, LY211795, LY 248908, mancozeb, maneb, mefenoxam, mepanipyrim, mepronil, metalaxyl, metalaxyl M, metconazole, metiram, metiram-zinc, metominostrobin, metrafenone, MON65500 (N-allyl-4,5-dimethyl-2-trimethylsilylthiophene-3- 15 carboxamide), myclobutanil, NTN0301, neoasozin, nickel dimethyldithiocarbamate, nitrothale-isopropyl, nuarimol, ofurace, organomercury compounds, orysastrobin, oxadixyl, oxasulfuron, oxolinic acid, oxpoconazole, oxycarboxin, pefurazoate, penconazole, pencycuron, phenazin oxide, phosphorus acids, phthalide, picoxystrobin, polyoxin D, polyram, probenazole, prochloraz, procymidone, propamocarb, propamocarb 20 hydrochloride, propiconazole, propineb, propionic acid, proquinazid, prothioconazole, pyraclostrobin, pyrazophos, pyrifenox, pyrimethanil, pyroquilon, pyroxyfur, pyrrolnitrin, quaternary ammonium compounds, quinomethionate, quinoxifen, quintozone, silthiofam (MON 65500), S-imazalil, simeconazole, sipconazole, sodium pentachlorophenate, spiroxamine, streptomycin, sulphur, tebuconazole, tecloftalam, tecnazene, tetraconazole, 25 thiabendazole, thifluzamide, 2-(thiocyanomethylthio)benzothiazole, thiophanate-methyl, thiram, tiadinil, timibenconazole, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triazbutil, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin A, vapam, vinclozolin, XRD-563, zineb, ziram, zoxamide and compounds of the formulae:



The compounds of formula (1) may be mixed with soil, peat or other rooting media for the protection of plants against seed-borne, soil-borne or foliar fungal diseases.

Some mixtures may comprise active ingredients which have significantly different physical, chemical or biological properties such that they do not easily lend themselves to the same conventional formulation type. In these circumstances other formulation types may be prepared. For example, where one active ingredient is a water insoluble solid and the other a water insoluble liquid, it may nevertheless be possible to disperse each active ingredient in the same continuous aqueous phase by dispersing the solid active ingredient as a suspension (using a preparation analogous to that of an SC) but dispersing the liquid active ingredient as an emulsion (using a preparation analogous to that of an EW). The resultant composition is a suspoemulsion (SE) formulation.

The invention is illustrated by the following Examples in which the following abbreviations are used:

ml = millilitres

g = grammes

ppm = parts per million

M⁺ = mass ion

s = singlet

d = doublet

br s = broad singlet

t = triplet

DMSO = dimethylsulphoxide

NMR = nuclear magnetic resonance

HPLC = high performance liquid chromatography

q = quartet

m = multiplet

ppm = parts per million

EXAMPLE 1

This Example illustrates the preparation of 2-(6-quinolinylloxy)-N-(4-methylpent-2-yn-4-yl) butyramide (Compound No. 2 of Table 1)

Stage 1: Preparation of 2-bromo-N-(4-methylpent-2-yn-4-yl) butyramide

Step 1: Preparation of 4-amino-4-methylpent-2-yne hydrochloride

3-Amino-3-methylbutyne (commercially available as 90% aqueous solution; 16.6g) was dissolved in dichloromethane (150ml), dried over sodium sulphate and

filtered to give a solution containing 14.9g of amine. To the stirred solution of amine under an atmosphere of nitrogen at ambient temperature was added dry triethylamine (48.4ml). 1,2-Bis-(chlorodimethylsilyl)ethane (38.98g) in dichloromethane (100ml) was then added dropwise, maintaining the reaction temperature at 15°C by cooling. The mixture was stirred for 3 hours, the colourless solid, which had formed during the reaction, was filtered from solution and the filtrate was evaporated under reduced pressure to give a paste. The paste was extracted into hexane and refiltered. The filtrate was evaporated under reduced pressure and the oil obtained was distilled to give 1-(1,1-dimethyl-2-propynyl)-2,2,5,5-tetramethyl-1-aza-2,5-disilacyclopentane, 21.5g, b.p. 41°C at 0.06 mm Hg pressure.

^1H NMR (CDCl_3) δ : 0.16(12H, s); 0.60(4H, s); 1.48(6H, s); 2.24(1H, s).

Step 2

The product from Step 1 (13.0g) in dry tetrahydrofuran (140ml) was cooled to -70°C under an atmosphere of nitrogen with stirring and a solution of *n*-butyl lithium (23.1ml of 2.5M solution in hexanes) was added at -65 to -70°C during 5 minutes. The mixture was allowed to warm to -5°C and methyl iodide (3.93ml) was added dropwise over 10 minutes. The reaction mixture was allowed to warm to 10°C when an exothermic reaction occurred. The mixture was maintained at 20°C by cooling for 2 hours then evaporated under reduced pressure to a small volume. The residue was dissolved in hexane, filtered to remove the insoluble material and evaporated under reduced pressure to give 1-(1,1-dimethyl-2-butynyl)-2,2,5,5-tetramethyl-1-aza-2,5-disilacyclopentane as a yellow oil, 13.0g.

^1H NMR (CDCl_3) δ : 0.10(12H, s); 0.56(4H, s); 1.40(6H, s); 1.72(3H, s).

Step 3

The product from Step 2 (13.0g) was added slowly to aqueous hydrochloric acid (35ml, 4M) at 0°C with stirring. The emulsion formed was stirred for 0.5 hours then taken to pH 14 with aqueous sodium hydroxide (4M) while maintaining the reaction mixture at 0°C by cooling in ice. The aqueous mixture was extracted into dichloromethane (three times) and the extracts combined, dried over sodium sulphate and filtered. The filtrate was made acidic by adding an excess of a saturated solution of hydrogen chloride in 1,4-dioxan. The mixture was concentrated under reduced pressure until a colourless precipitate was formed. Hexane was added to the suspension and the

solid was filtered from solution. The solid was washed with dry diethyl ether and placed under vacuum to remove any residual solvents to give the required product as a colourless solid, 5.0g.

^1H NMR (d_6 -DMSO) δ : 1.74(6H, s); 1.82(3H, s); 8.74(3H, br s).

5 Step 4: The preparation of 2-bromo-*N*-(4-methylpent-2-yn-4-yl) butyramide

The product from Step3 (5.0g) was dissolved in dry dichloromethane (200ml), cooled to 3°C with stirring then 2-bromobutyryl bromide (6.25g) was added followed by dropwise addition of dry triethylamine (10.93ml), maintaining the reaction at 5°C. The suspension, which had formed during the reaction, was stirred at ambient temperature for 1 hour then
10 water was added. The organic phase was separated, washed with water, dried over magnesium sulphate then evaporated under reduced pressure. The residue was fractionated by chromatography (silica; hexane / diethyl ether, 3:1 by volume) to give the required product, 5.2g, as a colourless solid, mp 79-81°C.

^1H NMR (CDCl_3) δ : 1.04(3H, t); 1.64(6H, s); 1.84(3H, s); 2.04-2.18(2H, m); 4.20-
15 4.24(1H, m); 6.46(1H, br s).

Stage 2

6-Hydroxyquinoline (0.46g) in dry *N,N*-dimethylformamide (10ml) was added dropwise to a stirred suspension of sodium hydride (0.10g, 80% dispersion in mineral oil) in dry *N,N*-dimethylformamide under an atmosphere of nitrogen at ambient temperature.

20 The green solution was stirred at ambient temperature for 1 hour and a solution of 2-bromo-*N*-(4-methylpent-2-yn-4-yl) butyramide (0.74g) in dry *N,N*-dimethylformamide (10ml) was added. The mixture was stirred at ambient temperature for 18 hours, poured into water, and extracted into diethyl ether (three times). The organic extracts were combined, washed with dilute aqueous sodium hydroxide, water (twice), dried over
25 magnesium sulphate and evaporated under reduced pressure to give a gum. The gum was fractionated by chromatography (silica; diethyl ether) to give the required product, 0.44g, as a colourless gum.

^1H NMR (CDCl_3) δ : 1.06-1.10(3H, t); 1.56-1.60(6H, d); 1.76(3H, s); 1.98-2.10(2H, m);
4.54-4.58(1H, m); 6.42(1H, s); 7.10(1H, m); 7.36-7.44(2H, m); 8.02-8.06(2H, m);
30 8.82(1H, d).

EXAMPLE 2

This Example illustrates the preparation of 2-(6-quinolinylloxy)-*N*-(1-*tert*-butyldimethylsilyloxy-4-methylpent-2-yn-4-yl)butyramide (Compound No. 50 of Table 1)

Stage 1: Preparation of 4-amino-1-hydroxy-4-methylpent-2-yne hydrochloride.

5 Step 1

1-(1,1-Dimethyl-2-propynyl)-2,2,5,5-tetramethyl-1-aza-2,5-disilacyclopentane (22.6g) in dry tetrahydrofuran (250ml) was cooled to -50°C under an atmosphere of nitrogen with stirring and a solution of *n*-butyl lithium (44ml, 2.5M solution in hexanes) was added dropwise over 10 minutes. The mixture was stirred for 0.5 hour, allowed to warm to -20°C then formaldehyde gas was bubbled through the mixture until no starting material remained, as determined by glc analysis. On complete reaction the mixture was treated with water, the ether phase separated and the aqueous phase was extracted with ethyl acetate (twice). The organic extracts were combined, washed with water (three times), dried over magnesium sulphate and evaporated under reduced pressure to give the required product (24.96g) as a pale yellow liquid.

^1H NMR (CDCl_3) δ : 0.00(12H, s); 0.46(4H, s); 1.32(6H, s); 4.10(2H, s).

Step 2

The product from Step 1 (24.96g) was treated with dilute aqueous hydrochloric acid (300ml) and stirred at ambient temperature for 0.5 hour. The mixture was washed with diethyl ether (twice), the aqueous phase was evaporated under reduced pressure, distilled with toluene (twice) to remove residual water and the residual solid obtained was triturated with hexane to give 4-amino-1-hydroxy-4-methylpent-2-yne hydrochloride (13.1g) as a cream coloured solid.

^1H NMR (CDCl_3) δ : 1.48(6H, s); 4.06(2H, s); 5.32(1H, s); 8.64(3H, s).

25 Stage 2: Preparation of 4-amino-1-*tert*-.butyldimethylsilyloxy-4-methylpent-2-yne.

4-Amino-1-hydroxy-4-methylpent-2-yne hydrochloride (4.40g) was dissolved in dry *N,N*-dimethylformamide (100ml) and triethylamine (4.44ml) was added. The suspension was stirred at ambient temperature for 10 minutes, imidazole (4.93g) was added followed by *t*-butyldimethylsilyl chloride (5.24g) in dry *N,N*-dimethylformamide (40ml). The mixture was stirred at ambient temperature for 18 hours then diluted with water. The mixture was extracted with diethyl ether (three times) and the organic extracts

were combined, washed with water (twice) then dried over magnesium sulphate and evaporated under reduced pressure to give the required product (6.88g) as a yellow liquid.

^1H NMR (CDCl_3) δ : 0.04(6H, s); 0.84(9H, s); 1.30(6H, s); 4.22(2H, s).

Stage 3: Preparation of 2-(6-quinolinyloxy)butyric acid.

5 Step 1: Preparation of methyl 2-(6-quinolinyloxy)butyrate

6-Hydroxyquinoline (25.1g), methyl 2-bromobutyrate (32.3g) and anhydrous potassium carbonate (23.0g) in dry *N,N*-dimethylformamide (100ml) were stirred at 100°C for 3 hours then stored at ambient temperature for 18 hours. The mixture was added to water, extracted with ethyl acetate (three times) and the extracts combined, washed with water (four times) then dried over magnesium sulphate. The solvent was evaporated under reduced pressure to give the required product (39.0g) as a red oil.

^1H NMR (CDCl_3) δ : 1.10-1.14(3H, t); 2.04-2.12(2H, q); 3.78(3H, s); 4.72-4.76(1H, t); 7.00(1H, d); 7.36-7.40(1H, m); 7.42-7.46(1H, m); 8.02-8.04(2H, d); 8.80(1H, d).

Step 2: Preparation of 2-(6-quinolinyloxy)butyric acid.

15 The product from Step 1 (38.8g) was stirred in a solution of sodium hydroxide (12.6g) in water (100ml) and heated to 90°C for 3 hours then cooled to ambient temperature. The solution was diluted with water and the aqueous phase was washed with ethyl acetate (twice), acidified to pH6 with aqueous hydrochloric acid then extracted with ethyl acetate (three times). The extracts were combined, dried over magnesium sulphate and the solvent evaporated under reduced pressure and the residue was washed with hexane to give the required product, 8.1g, as a yellow brown solid. The aqueous phase was re-extracted with ethyl acetate and processed as before to provide further required product (3.7g).

Stage 4

25 2-(6-Quinolinyloxy)butyric acid (0.61g), 4-amino-1-*tert*-butyldimethylsilyloxy-4-methylpent-2-yne (0.57g) and 4-dimethylaminopyridine (0.010g) in dry dichloromethane (10ml) were stirred and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (0.53g) was added. The mixture was stirred at ambient temperature for 3.5 hours, stored for 2 days, diluted with dichloromethane, washed with saturated aqueous sodium hydrogen carbonate (twice) and then with water. The organic phase was dried over magnesium sulphate and evaporated under reduced pressure to give a yellow oil. The oil was fractionated by chromatography (silica; hexane/ethyl acetate, 3:1 by volume) to give

30

a gum that was triturated with hexane to give the required product (0.12g) as a colourless solid, m.p.78-80°C.

¹H NMR (CDCl₃) δ: 0.08(6H, s); 0.88(9H, s); 1.06-1.10(3H, t); 1.62-1.66(6H, d); 2.00-2.10(2H, m); 4.28(2H, s); 4.54-4.58(1H, t); 6.44(1H, s); 7.10(1H, s); 7.36-7.44(2H, m);
5 8.02-8.06(2H, d); 8.80(1H, m).

The hexane washings were evaporated under reduced pressure to give an oil (0.53g) containing further required product.

EXAMPLE 3

This Example illustrates the preparation of 2-(6-quinolinyloxy)-*N*-(1-hydroxy-4-methylpent-2-yn-4-yl) butyramide (Compound No. 10 of Table1)

2-(6-Quinolinyloxy)-*N*-(1-tert.butyltrimethylsilyloxy-4-methylpent-2-yn-4-yl)butyramide (0.58g) in tetrahydrofuran (10ml) was stirred at 3-5°C and a solution of tetra *n*-butylammonium fluoride (2.64ml of 1M solution in tetrahydrofuran) was added dropwise over 5 minutes. On completion of addition, the mixture was stirred for 0.5 hour
15 at 0°C, 0.75 hour at ambient temperature then stored for 18 hours. The solvent was evaporated under reduced pressure and the residue portioned between ethyl acetate and aqueous ammonium chloride. The organic phase was separated, washed with aqueous ammonium chloride, brine, dried over magnesium sulphate and evaporated under reduced pressure to give a gum that was fractionated by chromatography (silica; hexane/ethyl
20 acetate, 1:2 by volume) to give the required product as a colourless glass (0.30g).

¹H NMR (CDCl₃) δ: 1.06-1.10(3H, t); 1.58(3H, s); 1.60(3H, s); 1.98-2.08(3H, m); 4.22-4.24(2H, d); 4.56-4.60(1H, t); 6.42(1H, s); 7.10(1H, s); 7.36-7.44(2H, m); 8.04-8.08(2H, d); 8.80(1H, m).

EXAMPLE 4

25 This Example illustrates the preparation of 2-(6-quinolinyloxy)-*N*-(1-methoxy-4-methylpent-2-yn-4-yl)butyramide (Compound No. 12 of Table 1)

Step 1: Preparation of 4-amino-1-methoxy-4 methylpent-2-yne hydrochloride.

To a stirred suspension of sodium hydride (0.45g, 80% dispersion in mineral oil) in dry *N,N*-dimethylformamide (2ml) under an atmosphere of nitrogen at ambient temperature
30 was added dropwise over 5 minutes a solution of 4-amino-1-hydroxy-4 methylpent-2-yne hydrochloride (0.75g) in dry *N,N*-dimethylformamide (20ml). The mixture was stirred for 2.75 hours at ambient temperature then a solution of methyl iodide (0.78g) in *N,N*-di-

methylformamide (5ml) was added. The reaction was stirred for 2.5 hours, stored for 18 hours then poured into water, extracted with diethyl ether (three times) and the organic extracts were combined. The combined organic phase was extracted with dilute hydrochloric acid (three times) and the aqueous acidic extracts were combined and
5 evaporated under reduced pressure. The residual solid was dried by evaporating under reduced pressure with toluene (twice) to give a yellow gum (0.8g) containing the required product. The compound was characterised from its NMR spectrum.

^1H NMR (CDCl_3) δ : 1.78(6H, s); 3.40(3H, s); 4.12(2H, s); 8.90(3H, br s).

Step 2

10 Triethylamine (0.54ml) was added to a stirred solution of the product from Step 1 (0.8g) in dry *N,N*-dimethylformamide (10ml). The solution was stirred for 5 minutes then 1-hydroxybenzotriazole (0.39g) and *N*-(3-dimethylaminopropyl)-*N'*-ethyl carbodiimide hydrochloride (0.55g) in dry *N,N*-dimethylformamide (5ml) were added. The mixture was stirred at ambient temperature for 2 hours, poured into water and the aqueous phase
15 extracted with ethyl acetate (three times). The organic extracts were combined, washed with water (three times), dried over magnesium sulphate and evaporated under reduced pressure to give a red oil. The oil was fractionated by chromatography (silica; ethyl acetate) to give the required product (0.15g) as a pale yellow gum.

^1H NMR (CDCl_3) δ : 1.06-1.12(3H, t); 1.62(6H, s); 2.00-2.08(2H, m); 3.32(3H, s); 4.08-
20 4.24(2H, s); 4.58-4.62(1H, t); 6.44(1H, s); 7.10(1H, m); 7.36-7.46(2H, m); 8.04-8.08(2H, d); 8.82(1H, m).

EXAMPLE 5

This Example illustrates the preparation of 2-(6-quinolinylloxy)-2-(ethoxy)-*N*-(2-methylpent-3-yn-2-yl) acetamide (Compound No. 2 of Table 17)

Step 1: Preparation of ethyl 2-(6-quinolinylloxy)-2-(ethoxy)acetate.

25 Potassium t-butoxide (3.15g) was dissolved in t-butyl alcohol (20ml), and stirred for 10 minutes at ambient temperature. 6-Hydroxyquinoline (3.0g) was added, the resulting dark green solution was stirred for 15 minutes, and then ethyl 2-chloro-2-ethoxyacetate (4.22g, 90% pure) added, followed by a catalytic amount of potassium iodide. The mixture was
30 stirred for 18 hours, poured into water and extracted with chloroform. The organic fraction was washed with brine and then water, and dried over magnesium sulphate. The solvent was evaporated to give a brown oil, which was purified by flash chromatography

on silica gel, eluting with a gradient of ethyl acetate:hexane from 1:2 to 4:1. Ethyl 2-(6-quinolinyloxy)-2-(ethoxy)acetate was obtained as a pale brown oil (3.68g).

^1H NMR (CDCl_3) δ ppm: 1.28 (3H, t); 1.31 (3H, t); 3.79 (1H, m); 3.90 (1H, m); 4.31 (2H, q); 5.68 (1H, s); 7.38 (2H, dd); 7.51 (1H, dd); 8.06 (2H, dd); 8.32 (1H, dd).

5 Step 2: Preparation of 2-(6-quinolinyloxy)-2-(ethoxy)acetic acid.

Ethyl 2-(6-quinolinyloxy)-2-(ethoxy)acetate (3.68g) was added to a solution of sodium hydroxide (0.589g) in water (10ml) and methanol (30ml), and stirred for 5 minutes. The solution was evaporated under reduced pressure, water added and the aqueous phase was washed with ethyl acetate. The aqueous phase was then acidified with hydrochloric acid and extracted with ethyl acetate. The extracts were combined, dried over magnesium sulphate and evaporated under reduced pressure to give 2-(6-quinolinyloxy)-2-(ethoxy)acetic acid, (1.47g), as a cream solid.

10 ^1H NMR (CDCl_3) δ ppm: 1.33 (3H, t); 3.98 (1H, m); 5.71 (1H, s); 7.44 (1H, dd); 7.52 (2H, m); 8.05 (1H, d); 8.20 (1H, d); 8.84 (1H, dd).

15 Step 3

Triethylamine (0.3ml) was added to a stirred solution of 4-amino-4-methylpent-2-yne hydrochloride (0.054g) in *N,N*-dimethylformamide (2 ml) giving a white suspension. 2-(6-Quinolinyloxy)-2-(ethoxy)acetic acid (0.1g) was then added followed by *N*-(3-dimethylaminopropyl)-*N'*-ethyl carbodiimide hydrochloride (0.077g) and a catalytic amount of 1-hydroxybenzotriazole, and the white suspension was stirred at ambient temperature for 18 hours. Water was added and the aqueous phase was extracted with ethyl acetate. The organic phase was washed with water, saturated sodium bicarbonate and brine, dried over magnesium sulphate and then evaporated under reduced pressure to give a yellow oil (0.117 g). This was purified by flash chromatography (silica gel; ethyl acetate/hexane 1:1 by volume) to give the required product (0.096g) as a colourless oil.

25 ^1H NMR (CDCl_3) δ ppm: 1.28 (3H, t); 1.64 (3H, s); 1.66 (3H, s); 1.80 (3H, s); 3.70 (1H, m); 3.88 (1H, m); 5.48 (1H, s); 6.79 (1H, bs); 7.37 (1H, dd); 7.49 (1H, d); 7.52 (1H, dd); 8.05 (1H, d); 8.08 (1H, d); 8.82 (1H, dd).

30

EXAMPLE 6

This Example illustrates the preparation of 2-(6-quinolinyloxy)-*N*-(2-methylpent-3-yn-2-yl)-3-methoxypropionamide (Compound No. 2 of Table 11)

Stage 1: Preparation of 2-bromo-*N*-(4-methylpent-2-yn-4-yl) 3-methoxypropionamide.

Step 1: Preparation of methyl 2-bromo-3-methoxypropionate

Methyl 2,3-dibromopropionate (21.9g) and trimethylamine *N*-oxide (0.1g) in methanol (8ml) were cooled to -5°C with stirring under an atmosphere of nitrogen. A methanolic solution of sodium methoxide, freshly prepared from sodium (2.25g) and methanol (24ml), was added dropwise over 15 minutes to the mixture, which was maintained below 0°C by cooling. On complete addition the mixture was stirred for a further 30 minutes then acetic acid (1ml) was added followed by diethyl ether (100ml). The mixture was filtered to remove insoluble salts and the filtrate evaporated under reduced pressure to give an oil, which was re-dissolved in a small volume of diethyl ether and re-filtered. The filtrate was evaporated under reduced pressure to give the required product (17.4g) as a pale yellow oil.

$^1\text{H NMR}$ (CDCl_3) δ : 3.41(3H, s); 3.74(1H, dd); 3.82(3H, s); 3.92(1H, dd); 4.34(1H, dd).

Step 2: Preparation of 2-bromo-3-methoxypropionic acid.

Methyl 2-bromo-3-methoxypropionate (1.00g) in tetrahydrofuran (8ml) was stirred at 10°C and lithium hydroxide monohydrate (0.21g) in water (1.5ml) was added dropwise. On completion of addition, the mixture was stirred for 1.5 hours, the colourless solution was evaporated under reduced pressure to a small volume then the aqueous solution was taken to pH 3 with dilute sulphuric acid. The mixture was extracted with diethyl ether (50ml) and the organic phase separated, washed with brine, dried over magnesium sulphate then evaporated under reduced pressure to give the required product (0.6g) as a colourless liquid.

$^1\text{H NMR}$ (CDCl_3) δ : 3.45(3H, s); 3.78(1H, m); 3.92(1H, m); 4.38(1H, m); 6.65(1H, br s).

Step 3: Preparation of 2-bromo-*N*-(4-methylpent-2-yn-4-yl) 3-methoxypropionamide.

2-Bromo-3-methoxypropionic acid (0.366g) was dissolved in dry dichloromethane (4ml) containing dry *N,N*-dimethylformamide (0.05ml) with stirring and oxalyl chloride (0.254g) was added. The mixture was stirred at ambient temperature for 2 hours then evaporated under reduced pressure to give 2-bromo-3-methoxypropionic acid chloride ($\text{C}=\text{O}$, ν 1780cm^{-1}). The acid chloride was dissolved in dry dichloromethane (6ml) and 4-amino-4-methylpent-2-yne hydrochloride (0.267g) was added then the mixture was cooled to 3°C and triethylamine (0.404g) was added dropwise, keeping the reaction temperature between $0-5^{\circ}\text{C}$. The suspension that had formed was stirred at

ambient temperature for 1 hour, diluted with further dichloromethane, washed with hydrochloric acid (2M) and the organic phase separated, dried over magnesium sulfate then evaporated under reduced pressure to give a gum. The gum was fractionated by chromatography (silica: hexane/ethyl acetate, 3:2 by volume) to give the required product (0.3g) as a colourless solid.

¹H NMR (CDCl₃) δ: 1.63(6H, s); 1.82(3H, s); 3.44(3H, s); 3.88(2H, m); 4.32(1H, m); 6.62(1H, s).

Stage 2

6-Hydroxyquinoline (0.083g), anhydrous potassium carbonate (0.087g) and 2-bromo-*N*-(4-methylpent-2-yn-4-yl) 3-methoxypropionamide (0.150g) in dry *N,N*-dimethylformamide (3ml) were stirred and heated to 80°C for 5 hours then stored for 72 hours at ambient temperature. The yellow suspension was diluted with water, extracted with ethyl acetate and the organic phase was separated then washed with water, dried over magnesium sulphate and evaporated under reduced pressure to a gum. The gum was fractionated by chromatography (silica; ethyl acetate: hexane, 3:2 by volume) to give the required product (0.055g) as a colourless gum.

¹H NMR (CDCl₃) δ: 1.58(3H, s); 1.60(3H, s); 1.78(3H, s); 3.46(3H, s); 3.95(2H, m); 4.78(1H, m); 6.60(1H, s); 7.18(1H, m); 7.40(1H, m); 7.50(1H, m); 8.08(2H, m); 8.83(1H, m).

20

EXAMPLE 7

This Example illustrates the preparation of 2-(6-quinazolinooxy)-*N*-(4-methylpent-2-yn-4-yl) butyramide (Compound No.2 of Table 20)

6-Hydroxyquinazoline (0.060g, prepared as described in *J. Chem. Soc.* (1952), 4985) and 2-bromo-*N*-(4-methylpent-2-yn-4-yl) butyramide (0.101g) were dissolved in dry *N,N*-dimethylformamide (2ml) containing anhydrous potassium carbonate (0.088g). The mixture was stirred and heated to 80°C for 5 hours then allowed to cool to ambient temperature and stored for 18 hours. The brown suspension was diluted with water, extracted into ethyl acetate and the organic phase separated, washed with water, dried over magnesium sulphate and evaporated under reduced pressure to give a pale brown oil. The oil was fractionated by chromatography (silica; ethyl acetate/hexane, 4:1 by volume) to give the title product (0.12g) as a pale brown gum.

¹H NMR (CDCl₃) δ: 1.04(3H, t); 1.53(6H, s); 1.72(3H, s); 1.95-2.05(2H, m); 4.54(1H, m); 6.30(1H, s); 7.14(1H, m); 7.60(1H, dd); 7.97(1H, d); 9.20(1H, s); 9.25(1H, s).

EXAMPLE 8

In a similar procedure to Example 1, 7-hydroxyisoquinoline (commercially available) and 2-bromo-*N*-(4-methylpent-2-yn-4-yl) butyramide were reacted to give 2-(7-isoquinolin-5 yloxy)-*N*-(4-methylpent-2-yn-4-yl) butyramide (Compound No. 2 of Table 39) as a colourless solid, m.p. 149-150°C.

¹H NMR (CDCl₃) δ: 1.08(3H, t); 1.58(3H, s); 1.59(3H, s); 1.77(3H, s); 1.98-2.12(2H, m); 4.58(1H, m); 6.40(1H, s); 7.26(1H, d); 7.42(1H, dd); 7.61(1H, d); 7.79(1H, d); 8.45(1H, 10 d); 9.14(1H, s).

EXAMPLE 9

In a similar procedure to Example 6, 7-hydroxyisoquinoline (commercially available) and 2-bromo-*N*-(4-methylpent-2-yn-4-yl) 3-methoxypropionamide were reacted to give 2-(7-isoquinolin-15 yloxy)-*N*-(2-methylpent-3-yn-2-yl)-3-methoxypropionamide (Compound No. 2 of Table 49) as a colourless solid, m.p. 155-156°C.

¹H NMR (CDCl₃) δ: 1.59(3H, s); 1.60(3H, s); 1.77(3H, s); 3.45(3H, s); 3.88-3.96(2H, m); 4.78(1H, m); 6.54(1H, s); 7.31(1H, d); 7.47(1H, dd); 7.61(1H, d); 7.80(1H, d); 8.46(1H, d); 9.14(1H, s).

EXAMPLE 10

20 This Example illustrates the fungicidal properties of compounds of formula (1).

The compounds were tested in a leaf disk assay, with methods described below.

The test compounds were dissolved in DMSO and diluted into water to 200 ppm. In the case of the test on *Pythium ultimum*, they were dissolved in DMSO and diluted into water to 20 ppm.

25 *Erysiphe graminis f.sp. hordei* (barley powdery mildew): Barley leaf segments were placed on agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed four days after inoculation as preventive fungicidal activity.

30 *Erysiphe graminis f.sp. tritici* (wheat powdery mildew): Wheat leaf segments were placed on agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated

with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed four days after inoculation as preventive fungicidal activity.

Puccinia recondita f.sp. tritici (wheat brown rust): Wheat leaf segments were placed on agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed nine days after inoculation as preventive fungicidal activity.

Septoria nodorum (wheat glume blotch): Wheat leaf segments were placed on agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed four days after inoculation as preventive fungicidal activity.

Pyrenophora teres (barley net blotch): Barley leaf segments were placed on agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed four days after inoculation as preventive fungicidal activity.

Pyricularia oryzae (rice blast): Rice leaf segments were placed on agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed four days after inoculation as preventive fungicidal activity.

Botrytis cinerea (grey mould): Bean leaf disks were placed on agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed four days after inoculation as preventive fungicidal activity.

Phytophthora infestans (late blight of potato on tomato): Tomato leaf disks were placed on water agar in a 24-well plate and sprayed with a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed four days after inoculation as preventive fungicidal activity.

Plasmopara viticola (downy mildew of grapevine): Grapevine leaf disks were placed on agar in a 24-well plate and sprayed a solution of the test compound. After allowing to dry completely, for between 12 and 24 hours, the leaf disks were inoculated with a spore suspension of the fungus. After appropriate incubation the activity of a compound was assessed seven days after inoculation as preventive fungicidal activity.

Pythium ultimum (Damping off): Mycelial fragments of the fungus, prepared from a fresh liquid culture, were mixed into potato dextrose broth. A solution of the test compound in dimethyl sulphoxide was diluted with water to 20ppm then placed into a 96-well microtiter plate and the nutrient broth containing the fungal spores was added. The test plate was incubated at 24°C and the inhibition of growth was determined photometrically after 48 hours.

The following Compounds [Compound No (Table)] gave at least 70% control of the following fungal infections at 200ppm:

Phytophthora infestans: 2(1), 10(1), 12(1), 50(1), 2(20).

15 *Plasmopara viticola*: 2(1).

Erysiphe graminis f.sp. hordei: 10(1), 2(17).


Erysiphe graminis f.sp. tritici: 10(1), 50(1),

The following Compounds gave at least 70% control of the following fungal infection at 20ppm:

20 *Pythium ultimum*: 2(1), 10(1), 12(1), 50(1), 2(11), 2(17), 2 (20).

CLAIMS

1. A compound of the general formula (1):



- 5 wherein one of X and Y is N and the other is CR or both of X and Y are N; R is H, halo, C₁₋₈ alkyl, C₃₋₆ cycloalkyl, C₂₋₈ alkenyl, C₂₋₈ alkynyl, C₁₋₈ alkoxy, C₁₋₈ alkylthio, nitro, amino, mono- or di-(C₁₋₆)alkylamino, mono- or di-(C₂₋₆)alkenyl-amino, mono- or di-(C₂₋₆)alkynylamino, formylamino, C₁₋₄ alkyl(formyl)amino, C₁₋₄ alkylcarbonylamino, C₁₋₄ alkyl(C₁₋₄ alkylcarbonyl)amino, cyano, formyl, C₁₋₄ alkylcarbonyl, C₁₋₄ alkoxycarbonyl, aminocarbonyl, mono- or di-(C₁₋₄)alkylamino-carbonyl, carboxy, C₁₋₄ alkylcarbonyloxy, aryl(C₁₋₄)alkylcarbonyloxy, C₁₋₄ alkyl-sulphinyl, C₁₋₄ alkylsulphonyl or C₁₋₄ alkylsulphonyloxy; R₁ is C₁₋₄ alkyl, C₂₋₄ alkenyl or C₂₋₄ alkynyl in which the alkyl, alkenyl and alkynyl groups are optionally substituted on their terminal carbon atom with one, two or three
- 10 halogen atoms, with a cyano group, with a C₁₋₄ alkylcarbonyl group, with a C₁₋₄ alkoxycarbonyl group or with a hydroxy group, or R₁ is alkoxyalkyl, alkylthioalkyl, alkylsulphinylalkyl or alkylsulphonylalkyl in which the total number of carbon atoms is 2 or 3, or R₁ is a straight-chain C₁₋₄ alkoxy group; R₂ is H, C₁₋₄ alkyl, C₁₋₄ alkoxymethyl or benzyloxymethyl in which the phenyl ring of the benzyl moiety is optionally substituted with C₁₋₄ alkoxy; R₃ and R₄ are
- 15 independently H, C₁₋₃ alkyl, C₂₋₃ alkenyl or C₂₋₃ alkynyl provided that both are not H and that when both are other than H their combined total of carbon atoms does not exceed 4, or R₃ and R₄ join with the carbon atom to which they are attached to form a 3 or 4 membered carbocyclic ring optionally containing one O, S or N
- 20 atom and optionally substituted with halo or C₁₋₄ alkyl; and R₅ is H, C₁₋₄ alkyl or C₃₋₆ cycloalkyl in which the alkyl or cycloalkyl group is optionally substituted with halo, hydroxy, C₁₋₆ alkoxy, C₁₋₆ alkylthio, cyano, C₁₋₄ alkylcarbonyloxy, aminocarbonyloxy or mono- or di(C₁₋₄)alkylaminocarbonyloxy, tri(C₁₋₄)alkyl-silyloxy, optionally substituted phenoxy, optionally substituted thienyloxy,
- 25

optionally substituted benzyloxy or optionally substituted thienylmethoxy, or R₅ is optionally substituted phenyl, optionally substituted thienyl or optionally substituted benzyl, in which the optionally substituted phenyl and thienyl rings of the R₅ values are optionally substituted with one, two or three substituents
5 selected from halo, hydroxy, mercapto, C₁₋₄ alkyl, C₂₋₄, alkenyl, C₂₋₄ alkynyl, C₁₋₄ alkoxy, C₂₋₄ alkenyloxy, C₂₋₄ alkynyloxy, halo (C₁₋₄)alkyl, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, halo(C₁₋₄)alkylthio, hydroxy(C₁₋₄)alkyl, C₁₋₄alkoxy(C₁₋₄)alkyl, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenoxy, benzyloxy, benzoyloxy, cyano, isocyano, thiocyanato, isothiocyanato, nitro, -NR'R", -NHCOR', -NHCONR'R",
10 -CONR'R", -SO₂R', -OSO₂R', -COR', -CR'=NR" or -N=CR'R", in which R' and R" are independently hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, C₁₋₄ alkoxy, halo(C₁₋₄)alkoxy, C₁₋₄ alkylthio, C₃₋₆ cycloalkyl, C₃₋₆ cycloalkyl(C₁₋₄)alkyl, phenyl or benzyl, the phenyl and benzyl groups being optionally substituted with halogen, C₁₋₄ alkyl or C₁₋₄ alkoxy.

15

2. A compound according to claim 1 wherein X is N and Y is CH or X and Y are both N.
3. A compound according to claim 1 or 2 wherein R₁ is methyl, ethyl, *n*-propyl,
20 2,2,2-trifluoromethyl, cyanomethyl, acetylmethyl, methoxycarbonylmethyl, methoxycarbonylethyl, hydroxymethyl, hydroxyethyl, methoxymethyl, methylthiomethyl, ethoxymethyl, 2-methoxyethyl, 2-methylthioethyl, methoxy, ethoxy, *n*-propoxy or *n*-butoxy.
- 25 4. A compound according to claim 1 or 2 wherein R₁ is ethyl, methoxy, ethoxy or methoxymethyl.
5. A compound according to any one of the preceding claims wherein R₂ is H.
- 30 6. A compound according to any one of the preceding claims wherein both R₃ and R₄ are methyl.

7. A compound according to any one of the preceding claims wherein R₅ is H, methyl, hydroxymethyl, methoxymethyl or 1-methoxyethyl.
8. A compound according to claim 1 wherein R₁ methyl, ethyl, *n*-propyl, 2,2,2-trifluoromethyl, cyanomethyl, acetylmethyl, methoxycarbonylmethyl, methoxycarbonylethyl, hydroxymethyl, hydroxyethyl, methoxymethyl, methylthiomethyl, ethoxymethyl, 2-methoxyethyl, 2-methylthioethyl, methoxy, ethoxy, *n*-propoxy or *n*-butoxy; R₂ is H; R₃ and R₄ are both methyl; and R₅ is H, methyl, hydroxymethyl, methoxymethyl or 1-methoxyethyl.
9. A process for preparing a compound according to claim 1 as herein described.
10. A fungicidal composition comprising a fungicidally effective amount of a compound of formula (1) and a suitable carrier or diluent therefor.
11. A method of combating or controlling phytopathogenic fungi which comprises applying a fungicidally effective amount of a compound of formula (1) as defined in claim 1 or a composition according to claim 10 to a plant, to a seed of a plant, to the locus of the plant or seed or to soil or any other plant growth medium.

PCT Application
GB0304631



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☒ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.